

## Product datasheet for **RC600031**

### **MET (NM\_001127500) Human Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	MET (NM_001127500) Human Tagged ORF Clone
Tag:	DDK-His
Symbol:	MET
Synonyms:	AUTS9; c-Met; DFNB97; HGFR; RCCP2
Mammalian Cell Selection:	None
Vector:	pCMV6-XL5-DDK-His (PS100068)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RC600031 representing leader sequence plus the extracellular domain region of NM\_001127500

Red=Cloning site Blue=ORF Green=Tags(s)

GTAATACGACTCACTATAGGGCGGCCGGAATTCGTGACTGGATCTGGTACCGAGGAGATCCGCCGCCG  
CGATCGCC

ATGAAGGCCCGCTGTGCTTGACCTGGCATCCTCGTGCTCCTGTTACCTTGGTGCAGAGGAGCAATG  
GGGAGTGTAAAGAGGCACTAGCAAAGTCCGAGATGAATGTGAATATGAAGTATCAGCTTCCCAACTTCAC  
CGCGGAAACACCCATCCAGAATGTCATTCTACATGAGCATCACATTTTCCTTGGTGCCACTAACTACATT  
TATGTTTTAAATGAGGAAGACCTTCAGAAGTTGCTGAGTACAAGACTGGGCCTGTGCTGGAACACCCAG  
ATTGTTTCCCATGTCAGGACTGCAGCAGCAAAGCAATTTATCAGGAGGTGTTTGGAAAGATAACATCAA  
CATGGCTCTAGTGTGACACCTACTATGATGATCAACTATTAGCTGTGGCAGCGTCAACAGAGGGACC  
TGCCAGCGACATGCTTTCCCAACAATCATACTGCTGACATACAGTCGGAGGTTCACTGCATATTCTCCC  
CACAGATAGAAGAGCCCAGCCAGTGTCTGACTGTGTGGTGAAGCGCCCTGGGAGCCAAAGTCTTTTCATC  
TGTAAAGGACCGTTTCATCAACTCTTTGTAGGCAATACCATAAAATCTTCTTATTTCCAGATCATCCA  
TTGCATTCGATATCAGTGAGAAGGCTAAAGGAAACGAAAGATGGTTTTATGTTTTGACGGACCAGTCTCT  
ACATTGATGTTTTACCTGAGTTCAGAGATTCTTACCCCATTAAGTATGTCCATGCCTTTGAAAGCAACAA  
TTTTATTTACTTCTTGACGGTCCAAAGGAAACTCTAGATGCTCAGACTTTTCACACAAGAATAATCAGG  
TTCTGTTCATAAACTCTGGATTGCATTCTACATGGAATGCCTCTGGAGTGTATTCTCACAGAAAAGA  
GAAAAAGAGATCCACAAAGAAGGAAGTGTAAATATACTTCAGGCTGCGTATGTGAGCAAGCCTGGGGC  
CCAGCTTGCTAGACAAATAGGAGCCAGCCTGAATGATGACATTTCTTTTCGGGTGTTCCGACAAAGCAAG  
CCAGATTCTGCCAACCATGGATCGATCTGCCATGTGTGATTCCTATCAAATATGTCAACGACTTCT  
TCAACAAGATCGTCAACAAAAACAATGTGAGATGTCTCCAGCATTTTTACGGACCAATCATGAGCACTG  
CTTTAATAGGACACTTCTGAGAAATTCATCAGGCTGTGAAGCGCCCGTATGAATATCGAACAGAGTTT  
ACCACAGCTTTGCAGCGGTTGACTTATTCATGGGTCAATTCAGCGAAGTCTCTAACATCTATATCCA  
CCTTCATTAAGGAGACCTCACCATAGCTAATCTTGGGACATCAGAGGGTCGCTTCATGCAGGTTGTGGT  
TTCTCGATCAGGACCATCAACCCCTCATGTGAATTTCTCTGGACTCCCATCCAGTGTCTCCAGAAGTG  
ATTGTGGAGCATACATTAACCAAAATGGCTACACACTGGTTATCACTGGGAAGAAGATCACGAAGATCC  
CATTGAATGGCTTGGGCTGCAGACATTTCCAGTCTGCAGTCAATGCCTCTGCCCCACCCTTTGTTC  
GTGTGGCTGGTGCACGACAAATGTGTGCGATCGGAGGAATGCCTGAGCGGGACATGGACTCAACAGATC  
TGCTGCCTGCAATCTACAAGTTTTCCCAATAGTGCACCCCTGAAGGAGGGACAAGGCTGACCATAT  
GTGGCTGGGACTTTGGATTTCCGAGGAATAATAAATTTGATTTAAGAAAACTAGAGTTCTCCTTGGAAA  
TGAGAGCTGCACCTTGAAGTGAAGGACAGATGAATACATTGAAATGCACAGTTGGTCTGCCATG  
AATAAGCATTTCAATATGTCCATAATTTCAAATGGCCACGGGACAACACAATACAGTACATTCTCT  
ATGTGGATCCTGTAATAACAAGTATTTCCGCGAAATACGGTCTATGGCTGGTGGCACTTACTTACTTT  
AACTGGAAATTACCTAACAGTGGGAATTTAGACACATTTCAATTGGTGGAAAAACATGTACTTTAAAA  
AGTGTGCAACAGTATTTCTGAATGTTATACCCAGCCCAAACCATTCAACTGAGTTTGTGTTAAAT  
TGAAAATTGACTTAGCCAACCGAGAGACAAGCATCTTCAGTTACCGTGAAGATCCCATTGTCTATGAAAT  
TCATCCAACCAAACTTTTTATTAGTACTTGGTGGAAAGAACCTCTCAACATTGTGAGTTTTCTATTTTGC  
TTTGCCAGTGGTGGGAGCACAATAACAGGTGTTGGGAAAAACCTGAATTCAGTTAGTGTCCCAGAAATGG  
TCATAAATGTGCATGAAGCAGGAAGGAACCTTACAGTGGCATGTCAACATCGCTCTAATTCAGAGATAAT  
CTGTTGTACCACTCCTTCCCTGCAACAGCTGAATCTGCAACTCCCCCTGAAAACCAAGCCTTTTTTCATG  
TTAGATGGGATCCTTTCCAAATACTTTGATCTCATTATGTACATAATCCTGTGTTTAAAGCCTTTTGAAA  
AGCCAGTGTGATCTCAATGGGCAATGAAAATGTACTGAAATTAAGGGAAATGATATTGACCCTGAAGC  
AGTTAAAGGTGAAGTGTAAAAGTTGAAATAAGAGCTGTGAGAAATATACTTACATTCTGAAGCCGTT  
TTATGCACGGTCCCAATGACCTGCTGAAATTAAGAGCGAGCTAAATATAGAGTGAAGCAAGCAATTT  
CTTCAACCGTCTTGGAAAAGTAATAGTTCAACCAGATCAGAATTTACA

ACGCGTTCAGGGCACTACAAGGATGACGACGATAAGGGATCTCATCATCACCATCACCATTAATGAGATC  
TGGTACCGATATCAAGCTTGTGACTCTAGA

**Protein Sequence:** >RC600031 representing signal peptide plus the extracellular domain region of NM\_001127500  
 Red=Cloning sites Green= DDK and 6XHIS Tags

```


MKAPAVLAPGILVLLFTLVQRSNGECKEALAKSEMNVNMKYQLPNFTAETPIQNVILHEHHIFLGATNYI
YVLNEEDLQKVAEYKTPVLEHPDCFCQDCSSKANLGGVWVDNINMALVVDYDDQLISCGSVNRGT
CQRHVFPNHHTADIQSEVHCIFSPQIEEPSQPCDCVVSALGAKVLSSVKDRFINFFVGTINSSYFPDHP
LHSISVRRLLKETKDFMFLTDQSYIDVLPFRDSYPIKYVHAFESNNFIYFLTVQRETLDAQTFHTRIIR
FCSINSLHSYMEMPLECILTEKRKRSTKKEVFNIIQAAYVSKPGAQLARQIGASLNDLILFGVFAQSK
PDSAEPMDRSAMCAFPKIKYVNDFFNKIVNKNVNRCLQHFYGNHEHCNRTLLRNSSGCEARRDEYRTEF
TTALQRVDLFGQFSEVLLTSISTFIKGLDTIANLGTSEGRFMQVVVSRSGPSTPHVNFLLDHPVSPEV
IVEHTLNQNGYTLVITGKKITKIPLNGLGCRHFQSCSQCLSAPPFVQCGWCHDKCVRSEECLSGTWTQOI
CLPAIYKVPNSAPLEGGTRLICGWDFGFRNNKFDLKKTRVLLGNESCTLTLSESTMNTLKCTVGPAM
NKHFNMSIIISNGHGTQYSTFSYVDPVITSISPKYGPAGGTLTLTGNYLNSGNSRHISIGGKCTLK
SVSNSILECYTPAQTIISTEFAVKLKIDLANRETSIFSYPREDPIVYEIHPKSFISTWWEKPLNIVSFLFC
FASGGSTITGVGNLNSVSVPRMIVNHEAGRNFVACQHRNSSEIICCTPSLQQLNLQLPLKTKAFFM
LDGILSKYFDLIYVHNPVFKPFKPVMI SMGNENVLEIKGNDIDPEAVKGEVLKVGKSCENIHLHSEAV
LCTVPNDLLKLNSELNIEWKQAISSSTVLGKIVIQPDQNT
  
```

TRSGTRSGDYKDDDDKGSHHHHHH

**Chromatograms:** [https://cdn.origene.com/chromatograms/mk8118\\_a04.zip](https://cdn.origene.com/chromatograms/mk8118_a04.zip)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_001127500

**ORF Size:** 2850 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the extra cellular domain of the protein with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001127500.1</a> , <a href="#">NP_001120972.1</a>
<b>RefSeq Size:</b>	6695 bp
<b>RefSeq ORF:</b>	4227 bp
<b>Locus ID:</b>	4233
<b>UniProt ID:</b>	<a href="#">P08581</a>
<b>Cytogenetics:</b>	7q31.2
<b>Protein Families:</b>	Druggable Genome, Protein Kinase, Transmembrane
<b>Protein Pathways:</b>	Adherens junction, Axon guidance, Colorectal cancer, Cytokine-cytokine receptor interaction, Endocytosis, Epithelial cell signaling in Helicobacter pylori infection, Focal adhesion, Melanoma, Pathways in cancer, Renal cell carcinoma
<b>MW:</b>	106.3 kDa
<b>Gene Summary:</b>	This gene encodes a member of the receptor tyrosine kinase family of proteins and the product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple human cancers. [provided by RefSeq, May 2016]