

## Product datasheet for **RG202784**

### IMPA1 (NM\_005536) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	IMPA1 (NM_005536) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	IMPA1
Synonyms:	IMP; IMPA; MRT59
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG202784 representing NM_005536 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCTGATCCTTGGCAGGAATGCATGGATTATGCAGTAACTCTAGCAAGACAAGCTGGAGAGGTAGTTT  
GTGAAGCTATAAAAAATGAAATGAATGTTATGCTGAAAAGTTCTCCAGTTGATTTGGTAACTGCTACGGA  
CCAAAAAGTTGAAAAATGCTTATCTCTCCATAAAGGAAAAGTATCCATCTCACAGTTTCATTGGTGAA  
GAATCTGTGGCAGCTGGGAAAAAAGTATCTTAACCGACAACCCACATGGATCATTGACCCATTGATG  
GAACAACAACTTTGTACATAGATTTCTTTTGTAGCTGTTTCAATTGGCTTTGCTGTAATAAAAAAGAT  
AGAATTTGGAGTTGTGTACAGTTGTGTGAAGGCAAGATGTACACTGCCAGAAAAGGAAAAGGTGCCTTT  
TGTAATGGTCAAAAACACAAGTCTCACAACAAGAAGATATTACCAAATCTCTCTTGGTGACTGAGTTGG  
GCTCTTCCAGAACACCAGAGACTGTGAGAATGGTTCTTTCTAATATGGAAAAGCTTTTTTGCATTCTGT  
TCATGGGATCCGGAGTGTGGAACAGCAGCTGTTAATAATGTGCCTTGTGGCAACTGGCGGAGCAGATGCA  
TATTATGAAATGGGAATTCAGTCTGGGATGTTGCAGGAGCTGGCATTATTGTTACTGAAGCTGGTGGCG  
TGCTAATGGATGTTACAGGTGGACCATTTGATTTGATGTCACGAAGAGTAATTGCTGCAAATAATAGAAT  
ATTAGCAGAAAAGGATAGCTAAAGAAATTCAGGTTATACCTTTGCAACGAGACGACGAAGAT

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA



[View online »](#)

**Protein Sequence:** >RG202784 representing NM\_005536  
 Red=Cloning site Green=Tags(s)

MADPWQECMDYAVTLARQAGEVVCEAIKNEMNMLKSSPVDLVTATDQKVEKMLISSIKEKYP SHSFIGE  
 ESVAAGEKSILTDNPTWIIDPIDGTTNFVHRFPFVAVSIGFAVNKKIEFGVYSCVEGKMYTARKGKGAF  
 CNGQKLQVSQQEDITKSLLVTELGSSRTPETVRMVL SNMEKLF CIPVHGIRSVGTA AVNMCLVATGGADA  
 YYEMGIHCWDVAGAGIIVTEAGGVLMDVTGGPFDLMSRRVIAANNRILAERIAKEIQVIPLQRDDED

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



**ACCN:** NM\_005536

**ORF Size:** 831 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_005536.2](#), [NP\\_005527.1](#)

**RefSeq Size:** 2349 bp

**RefSeq ORF:** 834 bp

**Locus ID:** 3612

**UniProt ID:** [P29218](#)

**Cytogenetics:** 8q21.13


**Domains:** inositol\_P

**Protein Families:** Druggable Genome

**Protein Pathways:** Inositol phosphate metabolism, Metabolic pathways, Phosphatidylinositol signaling system

**Gene Summary:** This gene encodes an enzyme that dephosphorylates myo-inositol monophosphate to generate free myo-inositol, a precursor of phosphatidylinositol, and is therefore an important modulator of intracellular signal transduction via the production of the second messengers myo-inositol 1,4,5-trisphosphate and diacylglycerol. This enzyme can also use myo-inositol-1,3-diphosphate, myo-inositol-1,4-diphosphate, scyllo-inositol-phosphate, glucose-1-phosphate, glucose-6-phosphate, fructose-1-phosphate, beta-glycerophosphate, and 2'-AMP as substrates. This enzyme shows magnesium-dependent phosphatase activity and is inhibited by therapeutic concentrations of lithium. Inhibition of inositol monophosphate hydrolysis and subsequent depletion of inositol for phosphatidylinositol synthesis may explain the anti-manic and anti-depressive effects of lithium administered to treat bipolar disorder. Alternative splicing results in multiple transcript variants encoding distinct isoforms. A pseudogene of this gene is also present on chromosome 8q21.13. [provided by RefSeq, Dec 2014]

Product images:



Circular map for RG202784