

Product datasheet for **RG222167**

Glucocorticoid Receptor (NR3C1) (NM_001018075) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Glucocorticoid Receptor (NR3C1) (NM_001018075) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	NR3C1
Synonyms:	GCCR; GCR; GCRST; GR; GRL
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RG222167 representing NM_001018075
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGACTCCAAGAATCATTAACTCCTGGTAGAGAAGAAAACCCAGCAGTGTGCTTGCTCAGGAGAGGG
 GAGATGTGATGGACTTCTATAAAACCTAAGAGGAGGAGCTACTGTGAAGTTTTCTGCGTCTTACCCTC
 ACTGGCTGTCGCTTCTCAATCAGACTCCAAGCAGCGAAGACTTTTGGTTGATTTTCCAAAAGGCTCAGTA
 AGCAATGCGCAGCAGCCAGATCTGTCCAAGCAGTTTCACTCTCAATGGGACTGTATATGGGAGAGACAG
 AAACAAAAGTGATGGAAATGACCTGGGATCCCACAGCAGGGCCAAATCAGCCTTCTCGGGGAAAC
 AGACTTAAAGCTTTTGAAGAAAGCATTGCAAACCTCAATAGGTCGACCAGTGTCCAGAGAACCCCAAG
 AGTTCAGCATCCACTGCTGTGTCTGCTGCCCCACAGAGAAGGAGTTTCCAAAACCTCACTCTGATGAT
 CTTCAGAACAGCAACATTTGAAGGGCCAGACTGGCACCAACGGTGGCAATGTGAAATTGTATACCACAGA
 CCAAAGCACCTTTGACATTTTGACAGATTGGAGTTTTCTTCTGGGTCCCAGGTAAAGAGACGAATGAG
 AGTCCTTGAGATCAGACCTGTTGATAGATGAAAACCTGTTTGCTTCTCCTCTGGCGGGAGAAGACGATT
 CATTCTTTTGAAGGAACTCGAATGAGGACTGCAAGCCTCTCATTACCAGACTAAACCCAAAAT
 TAAGGATAATGGAGATCTGGTTTTGTCAAGCCCAGTAATGTAACACTGCCCAAGTGAACACAGAAAA
 GAAGATTTTCATCGAAGCTGACCCCTGGGGTAATTAAGCAAGAGAACTGGGCACAGTTTACTGTCAGG
 CAAGCTTCTCGGAGCAATATAATTGGTAATAAAATGTCTGCCATTTCTGTTTATGGTGTGAGTACCTC
 TGGAGGACAGATGTACCACTATGACATGAATACAGCATCCCTTCTCAACAGCAGGATCAGAAGCCTATT
 TTTAATGTCATTCCACCAATCCCGTTGGTCCGAAAATGGAATAGGTGCCAAGGATCTGGAGATGACA
 ACTTGACTTCTCGGGACTCTGAACCTCCCTGGTCCGAAACAGTTTTTTCTAATGGCTATTCAAGCCCCAG
 CATGAGACCAGATGTAAGCTCTCCTCCATCCAGCTCCTCAACAGCAACAACAGGACCACCTCCCAAACCTC
 TGCTGGTGTGCTCTGATGAAGCTTCAAGATGTCATTATGGAGTCTTAACTTGTGGAAGCTGTAAGTTT
 TCTTCAAAGAGCAGTGAAGGACAGCACAATTACCTATGTGCTGGAAGGAATGATTGCATCATCGATAA
 AATTCGAAGAAAAAACTGCCAGCATGCCGCTATCGAAAATGTCTTCAAGGCTGGAATGAACCTGGAAGCT
 CGAAAAACAAAGAAAAAATAAAAGGAATTCAGCAGGCCACTACAGGAGTCTCACAAGAAACCTCTGAAA
 ATCCTGGTAACAAAACAATAGTTCTGCAACGTTACCACAACCTACCCCTACCCTGGTGTCACTGTTGGA
 GGTATTGAACCTGAAGTGTATATGCAGGATATGATAGCTCTGTTCCAGACTCAACTGGAGGATCATG
 ACTACGCTCAACATGTTAGGAGGGCGCAAGTATTGCAGCAGTGAATGGGCAAAGGCAATACCAGGTT
 TCAGGAACCTACACCTGGATGACCAATGACCCTACTGCAGTACTCCTGGATGTTTCTTATGGCATTTC
 TCTGGGTGGAGATCATATAGACAATCAAGTGCAAACCTGCTGTGTTTTGCTCCTGATCTGATTATTAAT
 GAGCAGAGAATGACTCTACCCTGCATGTACGACCAATGTAACACATGCTGTATGTTTCTCTGAGTTAC
 ACAGGCTTCAGGTATCTTATGAAGAGTATCTCTGTATGAAAACCTTACTGCTTCTCTTTCAGTTCTTAA
 GGACGGTCTGAAGAGCCAAGAGCTATTTGATGAAATTAGAATGACCTACATCAAAGAGCTAGGAAAAAGCC
 ATTGTCAGAGGGAAGGAACTCCAGCCAGAAGTGGCAGCGGTTTTATCAACTGACAAAACCTTTGGATT
 CTATGCATGAAGTGGTTGAAAATCTCCTTAACTATTGCTTCCAAACATTTTGGATAAGACCATGAGTAT
 TGAATTCCTCCGAGATGTTAGCTGAAATCATCACCATCAGATACCAAAATATTCAATGGAAATATCAAA
 AAACCTTCTGTTTCATCAAAG

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

Protein Sequence: >RG222167 representing NM_001018075
 Red=Cloning site Green=Tags(s)


MDSKESLTPGREENPSSVLAQERGDVMDFYKTLRGGATVKVSASSPSLAVASQSDSKQRLLVDFPKGSV
 SNAQQPDL SKAVSL SMGLYMGETETKVMGNDLGFPOQQQISLSSGETDLKLL EESIANLNRSTVSPENPK
 SSASTAVSAAPTEKEFPKTHSDVSSEQHLKGQGTGNGGNVKLYTTDQSTFDILQDLEFSSGSPGKETNE
 SPWRSDLLIDENCLL SPLAGEDDSFLL EGNSEDCKPLILPDTKPKIKDNGDLVLSPPSNVTLPQVKTEK
 EDFIELCTPGVIKQEKLGTVYCCQASFPGANIIGNKMSAISVHGVSTSGGQMYHYDMNTASLSQQDQKPI
 FNVIPPIPVGSENWNRCCQSGDDNLTSLGTLNFPGRTVFSNGYSSPSMRPDVSSPPSSSSTATTGPPPKL
 CLVCSDEASGCHYGVLTCGSCVKVFFKRAVEGQHNYLCAGRNDICIIDKIRRKNCPACRYRKLQAGMNLEA
 RKTKKKIKGIQQATTGVSQETSENPGNKTIVPATLPQLTPTLVSLLEVIEPEVLYAGYDSSVPDSTWRIM
 TTLNMLGGRQVIAAVKWAKAIPGFRNLHLDDQMTLLQYSWMFLMAFALGWRYSYRQSSANLLCFAPDLIIN
 EQRMTLPCMYDQCKHMLYVSELHRLQVSYEEYLCMKTL LLLSSVPKDGLKSQELFDEIRMTYIKELGKA
 IVKREGNSSQNWRFYQLTKLLDSMHEVVENLLNYCFQTFLDKTM SIEFPEMLAEIITNQIPKYSNGNIK
 KLLFHQK

TRTRPLE - GFP Tag - V

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:


ACCN: NM_001018075

ORF Size: 2331 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001018075.1](#), [NP_001018085.1](#)

RefSeq Size: 6517 bp

RefSeq ORF: 2334 bp

Locus ID: 2908

UniProt ID: [P04150](#)

Cytogenetics: 5q31.3

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Neuroactive ligand-receptor interaction

Gene Summary: This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175). [provided by RefSeq, Feb 2011]