

## Product datasheet for **RG222342**

### Glucocorticoid Receptor (NR3C1) (NM\_001018077) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Glucocorticoid Receptor (NR3C1) (NM_001018077) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Glucocorticoid Receptor
Synonyms:	GCCR; GCR; GCRST; GR; GRL
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>RG222342 representing NM\_001018077  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGACTCCAAGAATCATTAACTCCTGGTAGAGAAGAAAACCCAGCAGTGTGCTTGCTCAGGAGAGGG  
 GAGATGTGATGGACTTCTATAAAACCCTAAGAGGAGGAGCTACTGTGAAGTTTTCTGCGTCTTACCCTC  
 ACTGGCTGTCGCTTCTCAATCAGACTCCAAGCAGCGAAGACTTTTGGTTGATTTTCCAAAAGGCTCAGTA  
 AGCAATGCGCAGCAGCCAGATCTGTCCAAGCAGTTTCACTCTCAATGGGACTGTATATGGGAGAGACAG  
 AAACAAAAGTGATGGAAATGACCTGGGATTTCCACAGCAGGGCCAAATCAGCCTTCTCGGGGAAAC  
 AGACTTAAAGCTTTTGAAGAAAGCATTGCAAACCTCAATAGGTCGACCAGTGTCCAGAGAACCCCAAG  
 AGTTCAGCATCCACTGCTGTGTCTGCTGCCCCACAGAGAAGGAGTTTCCAAAACCTCACTCTGATGAT  
 CTTCAGAACAGCAACATTTGAAGGGCCAGACTGGCACCAACGGTGGCAATGTGAAATTGTATACCACAGA  
 CCAAAGCACCTTTGACATTTTGCAAGATTTGGAGTTTTCTTCTGGGTCCCAGGTAAAGAGACGAATGAG  
 AGTCCTTGAGATCAGACCTGTTGATAGATGAAAACCTGTTTGCTTCTCCTCTGGCGGGAGAAGACGATT  
 CATTCTTTTGAAGGAACTCGAATGAGGACTGCAAGCCTCTCATTACCAGACTAAACCCAAAAT  
 TAAGGATAATGGAGATCTGGTTTTGTCAAGCCCAGTAATGTAACACTGCCCAAGTGAACACAGAAAA  
 GAAGATTTTCATCGAACTCTGCACCCCTGGGGTAATTAAGCAAGAGAACTGGGCACAGTTTACTGTCAGG  
 CAAGCTTCTCGGAGCAATATAATTGGTAATAAAATGTCTGCCATTTCTGTTTATGGTGTGAGTACCTC  
 TGGAGGACAGATGTACCACTATGACATGAATACAGCATCCCTTCTCAACAGCAGGATCAGAAGCCTATT  
 TTTAATGTCATTCCACCAATCCCGTTGGTCCGAAAATGGAATAGGTGCCAAGGATCTGGAGATGACA  
 ACTTGACTTCTCTGGGGACTCTGAACCTCCCTGGTCCGAAACAGTTTTTTCTAATGGCTATTCAAGCCCCAG  
 CATGAGACCAGATGTAAGCTCTCCTCCATCCAGCTCCTCAACAGCAACAACAGGACCACCTCCCAAACCTC  
 TGCTGGTGTGCTCTGATGAAGTTCAGGATGTCATTATGGAGTCTTAACTTGTGGAAGCTGTAAGTTT  
 TCTTCAAAGAGCAGTGGAAAGGACAGCACAATTACCTATGTGCTGGAAGGAATGATTGCATCATCGATAA  
 AATTCGAAGAAAAAACTGCCAGCATGCCGCTATCGAAAATGTCTTCAAGGCTGGAATGAACCTGGAAGCT  
 CGAAAAACAAAGAAAAAATAAAAGGAATTCAGCAGGCCACTACAGGAGTCTCACAAGAAACCTCTGAAA  
 ATCCTGGTAACAAAACAATAGTTCCTGCAACGTTACCACAACCTACCCCTACCCTGGTGTCACTGTTGGA  
 GGTATTGAACCTGAAGTGTATATGCAGGATATGATAGCTCTGTTCCAGACTCAACTTGGAGGATCATG  
 ACTACGCTCAACATGTTAGGAGGGCGCAAGTATTGCAGCAGTAAAATGGGCAAAGGCAATACCAGGTT  
 TCAGGAACCTACACCTGGATGACCAATGACCCTACTGCAGTACTCCTGGATGTTTCTTATGGCATTTC  
 TCTGGGTGGAGATCATATAGACAATCAAGTGCAAACCTGCTGTGTTTTGCTCCTGATCTGATTATTAAT  
 GAGCAGAGAATGACTCTACCCTGCATGTACGACCAATGTAACACATGCTGTATGTTTCTCTGAGTTAC  
 ACAGGCTTCAGGTATCTTATGAAGAGTATCTCTGTATGAAAACCTTACTGCTTCTCTTTCAGTTCTTAA  
 GGACGGTCTGAAGAGCCAAGAGCTATTTGATGAAATTAGAATGACCTACATCAAAGAGCTAGGAAAAAGCC  
 ATTGTCAGAGGGAAGGAACTCCAGCCAGAAGTGGCAGCGGTTTTATCAACTGACAAAACCTTTGGATT  
 CTATGCATGAAGTGGTTGAAAATCTCCTTAACTATTGCTTCCAAACATTTTGGATAAGACCATGAGTAT  
 TGAATTCCTCCGAGATGTTAGCTGAAATCATCACCATCAGATACCAAAATATTCAATGGAAATATCAAA  
 AAACCTTCTGTTTCATCAAAG

**ACGCGT**ACGCGGCCGCTCGAG - GFP Tag - GTTTAA

**Protein Sequence:** >RG222342 representing NM\_001018077  
 Red=Cloning site Green=Tags(s)

```
MDSKESLTPGREENPSSVLAQERGDVMDFYKTLRGGATVKVSASSPSLAVASQSDSKQRRLLVDFPKGSV
SNAQQPDL SKAVSL SMGLYMGETETKVMGNLGFPOQQQISLSSGETDLKLL EESIANLNRSTVSPENPK
SSASTAVSAAPTEKEFPKTHSDVSSEQHLKGQGTNGGNVKLYTTDQSTFDILQDLEFSSGSPGKETNE
SPWRSDLLIDENCLL SPLAGEDDSFLL EGNSEDCKPLILPDTKPKIKDNGDLVLSPPSNVTL PQVKTEK
EDFIELCTPGVIKQEKLGTVYQCASFPGANIIGNKMSAISVHGVSTSGGQMYHYDMNTASLSQQQDQKPI
FNVIPPIPVGSENWNR CQSGDDNLTSLGTLNFPGRTVF SNGYSSPSMRPDVSSPPSSSSTATTGPPPKL
CLVCSDEASGCHYGVLTCG SCKVFFKRAVEGQHNYL CAGRNDCIIDKIRRKNCPACRYRKLQAGMNLEA
RKTKKKIKGIQQATTGVSQETSENPGNKTIVPATLPQLTPTLVSLLEVIEPEVLYAGYDSSVPDSTWRIM
TTLNMLGGRQVIAAVK WAKAIPGFRNLHLDQMTLLQYSWMFLMAFALGWRSYRQSSANLLCFAPDLIIN
EQRMTLPCMYDQCKHML YVSELHRLQVSYEEYLCMKTL LLLSSVPKDGLKSQELFDEIRMTYIKELGKA
IVKREGNSSQNWR FYQLTKLLDSMHEVVENLLNYCFQ TFLDKTMSIEFPEMLAEIITNQIPKYSNGNIK
KLLFHQK
```

TRTRPLE - GFP Tag - V

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_001018077


**ORF Size:** 2331 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001018077.1</a></u> , <u><a href="#">NP_001018087.1</a></u>
<b>RefSeq Size:</b>	7286 bp
<b>RefSeq ORF:</b>	2334 bp
<b>Locus ID:</b>	2908
<b>UniProt ID:</b>	<u><a href="#">P04150</a></u>
<b>Cytogenetics:</b>	5q31.3
<b>Protein Families:</b>	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors
<b>Protein Pathways:</b>	Neuroactive ligand-receptor interaction
<b>Gene Summary:</b>	This gene encodes glucocorticoid receptor, which can function both as a transcription factor that binds to glucocorticoid response elements in the promoters of glucocorticoid responsive genes to activate their transcription, and as a regulator of other transcription factors. This receptor is typically found in the cytoplasm, but upon ligand binding, is transported into the nucleus. It is involved in inflammatory responses, cellular proliferation, and differentiation in target tissues. Mutations in this gene are associated with generalized glucocorticoid resistance. Alternative splicing of this gene results in transcript variants encoding either the same or different isoforms. Additional isoforms resulting from the use of alternate in-frame translation initiation sites have also been described, and shown to be functional, displaying diverse cytoplasm-to-nucleus trafficking patterns and distinct transcriptional activities (PMID:15866175). [provided by RefSeq, Feb 2011]

Product images:



Circular map for RG222342