

## Product datasheet for **RG238199**

### Ikaros (IKZF1) (NM\_001291838) Human Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Ikaros (IKZF1) (NM_001291838) Human Tagged ORF Clone
Tag:	TurboGFP
Symbol:	IKZF1
Synonyms:	CVID13; Hs.54452; IK1; IKAROS; LyF-1; LYF1; PPP1R92; PRO0758; ZNFN1A1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)
ORF Nucleotide Sequence:	>RG238199 representing NM_001291838. Blue=ORF Red=Cloning site Green=Tag(s)

```
GCTCGTTTGTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGGATGCTGATGAGGGTCAAGACATGTCCCAAGTTTCAGGGAAGGAAAGCCCCCTGTAAGCGTACT
CCAGATGAGGGCGATGAGCCATGCCGATCCCCGAGGACCTCTCCACCACCTCGGGAGGACAGCAAAGC
TCCAAGAGTGACAGAGTCGTGGGAGAACGGCCCTTCCAGTCAATCAGTGCAGGGGCTCATTACCCAG
AAGGGCAACCTGCTCCGGCACATCAAGTGCATTCCGGGGAAGCCCTCAAATGCCACCTCTGCAAC
TACGCTGCCCGGAGGACGCCCTCACTGGCCACCTGAGGACGCACTCCGTTGGTAAACCTCACAAA
TGTGGATATTGTGGCCGAAGCTATAAACAGCGAAGCTCTTTAGAGGAACATAAAGAGCGCTGCCACAAC
TACTTGAAAGCATGGGCCTTCCGGGCACACTGTACCCAGTCAATTAAGAAGAACTAATCACAGTGAA
ATGGCAGAAGACCTGTGCAAGATAGGATCAGAGAGATCTCTCGTCTGGACAGACTAGCAAGTAACGTC
GCCAAACGTAAGAGCTCTATGCCTCAGAAATTTCTTGGGACAAGGGCCTGTCCGACACGCCCTACGAC
AGCAGCGCCAGCTACGAGAAGGAGAACGAAATGATGAAGTCCCACGTGATGGACCAAGCCATCAACAAC
GCCATCAACTACCTGGGGCCGAGTCCCTGCGCCCGTGGTGCAGACGCCCGGGCGGTTCCGAGGTG
GTCCCGGTATCAGCCGATGTACCAGTGCACAAGCCGCTCGCGGAGGGCACCCCGCGTCCAACCAC
TCGGCCAGGACAGCGCCGTGGAGAACCTGCTGCTGCTCCTCAAGGCCAAGTTGGTGCCTCGGAGCGC
GAGGCGTCCCCGAGCAACAGCTGCCAAGACTCCACGGACACCGAGAGCAACAACGAGGAGCAGCGCAGC
GGTCTCATCTACCTGACCAACCACATCGCCCCGACGCGCAACGGGCTGTGCTCAAGGAGGAGCAC
CGCGCCTACGACCTGCTGCGCGCCGCTCCGAGAAGTCCGAGGACGCGCTCCGCGTGGTACGACCCAGC
GGGGAGCAGATGAAGGTGTACAAGTGCGAACACTGCCGGTGCTCTTCTGGATCACGTATGTACACC
ATCCACATGGGCTGCCACGGCTCCGTGATCCTTTTGTGAGTCAACATGTGCGGCTACCACAGCCAGGAC
CGGTACGAGTTCTCGTCGCACATAACGCGAGGGGAGCACCGCTTCCACATGAGC
ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAAAC
```




View online »


**Protein Sequence:** >Peptide sequence encoded by RG238199  
 Blue=ORF Red=Cloning site Green=Tag(s)

MDADEGQDMSQVSGKESPPVSDTPDEGDEPMPPIPEDLSTTSGGQQSSKSDRVVGERPFQCNQCGASFTQ  
 KGNLLRHIKLSHSGEKPFKCHLCNYACRRRDALTGHLRTHSVGKPHKCGYCGRSYKQRSSLEEKRCRCHN  
 YLESMGLPGTLYPVIKEETNHSEMAEDLCKIGSERSLVDRLASNVAKRKSMPQKFLGDKGLSDTPYD  
 SSASYEKENEMMKSHVMDQAINNAINYLGAE SLRPLVQTPPGGSEVVPVISPMPQLHKPLAEGTPRSNH  
 SAQDSAVENLLLLSKAKLVPSEREASPSNSCQDSTDTE SNNEEQRSGLIYLTNHIAPHARNGLSLKEEH  
 RAYDLLRAASENSQDALRVVSTSGEQMKVYKCEHCRVFLDHVMTIHMGC HGRDPFECNMGYHSQD  
 RYEFSSHITRGEHRFHMS  
**TRTRPLEMESDESGLPAMEIECRITGTLNGVEFELVGGGEGTPEQGRMTNKMKSTKGALTFSPYLLSHV**  
 MGYGFYHFGTYPSTYENPFLHAINNGGYNTRIEKYEDGGVLHVSFSYRYEAGRVIGDFKVMGTGFPED  
 SVIFTDKIIRSNATVEHLHPMGDNDLDGSFTRTFSLRDGGYSSVVD SHMHFKSAIHPSILQNGGPMFA  
 FRRVEEDHSNTELGIVEYQHAFKTPDADAGEERV

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**Plasmid Map:**


**ACCN:** NM\_001291838

**ORF Size:** 1296 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**RefSeq:** [NM\\_001291838.2](#)

**RefSeq Size:** 6029 bp

**RefSeq ORF:** 1299 bp

**Locus ID:** 10320

**UniProt ID:** [Q13422](#)

**Cytogenetics:** 7p12.2

**Protein Families:** Druggable Genome, Transcription Factors

**MW:** 48.7 kDa

**Gene Summary:**

This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL). [provided by RefSeq, May 2014]