

Product datasheet for RR212804L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Abhd6 (NM_001007680) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Abhd6 (NM 001007680) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Abhd6

Synonyms: MGC94917

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001007680

ORF Size: 1011 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR212804).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001007680.1</u>, <u>NP 001007681.1</u>

 RefSeq Size:
 2225 bp

 RefSeq ORF:
 1014 bp

 Locus ID:
 305795

 UniProt ID:
 Q5XI64

 Cytogenetics:
 15p14







Gene Summary:

Lipase that preferentially hydrolysis medium-chain saturated monoacylglycerols including 2-arachidonoylglycerol (By similarity). Through 2-arachidonoylglycerol degradation may regulate endocannabinoid signaling pathways. Also has a lysophosphatidyl lipase activity with a preference for lysophosphatidylglycerol among other lysophospholipids (By similarity). Also able to degrade bis(monoacylglycero)phosphate (BMP) and constitutes the major enzyme for BMP catabolism. BMP, also known as lysobisphosphatidic acid, is enriched in late endosomes and lysosomes and plays a key role in the formation of intraluminal vesicles and in lipid sorting (By similarity).[UniProtKB/Swiss-Prot Function]