

Product datasheet for SC311387

BIN1 (NM_139349) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: BIN1 (NM_139349) Human Untagged Clone

Tag: Tag Free

Symbol: BIN1

Synonyms: AMPH2; AMPHL; CNM2; SH3P9

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Fully Sequenced ORF:

>SC311387 representing NM_139349.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGGATCGCC


ATGGCAGAGATGGGCAGTAAAGGGGTGACGGCGGGAAAGATCGCCAGCAACGTGCAGAAGAAGCTCACC CGCGCGCAGGAGAAGGTTCTCCAGAAGCTGGGGAAGGCAGATGAGACCAAGGATGAGCAGTTTGAGCAG GCCTCCGTCAAAGCCATGCACGAGGCTTCCAAGAAGCTGAATGAGTGTCTGCAGGAGGTGTATGAGCCC CAGAAGCTGGTGGACCAGGCGCTGCTGACCATGGACACGTACCTGGGCCAGTTCCCCGACATCAAGTCA CGCATTGCCAAGCGGGGGCGCAAGCTGGTGGACTACGACAGTGCCCGGCACCACTACGAGTCCCTTCAA ACTGCCAAAAAGAAGGATGAAGCCAAAATTGCCAAGGCCGAGGAGGAGCTCATCAAAGCCCAGAAGGTG TTTGAGGAGATGAATGTGGATCTGCAGGAGGAGCTGCCGTCCCTGTGGAACAGCCGCGTAGGTTTCTAC GTCAACACGTTCCAGAGCATCGCGGGCCTGGAGGAAAACTTCCACAAGGAGATGAGCAAGCTCAACCAG AACCTCAATGATGTGCTGGTCGGCCTGGAGAAGCAACACGGGAGCAACACCTTCACGGTCAAGGCCCAG CCCAGTGACAACGCGCCTGCAAAAGGGAACAAGAGCCCTTCGCCTCCAGATGGCTCCCCTGCCGCCACC CCCGAGATCAGAGTCAACCACGAGCCAGAGCCGGCCGGGGGCCACGCCCGGGGCCACCCTCCCCAAG TCCCCATCTCAGCCCACAGAGAGTCCAGCCGGCAGCCTTCCGGGGGAGCCCAGCGCTGCCGAGGGC ACCTTTGCTGTGCCCTGGCCCAGCCAGACGCCGAGCCGGGGCCTGCCCAACCAGCAGAGGCCTCGGAG GTGGCGGGTGGGACCCAACCTGCGGCTGGAGCCCAGGAGCCAGGGGAAGCGGCGCAAGTGAAGCAGCC TCCAGCTCTCTTCCTGCTGTCGTGGAGACCTTCCCAGCAACTGTGAATGGCACCGTGGAGGGCGGC AGTGGGGCCGGGCCTTGGACCTGCCCCCAGGTTTCATGTTCAAGGTACAGGCCCAGCACGACTACACG GCCACTGACACAGACGAGCTGCAGCTCAAGGCTGGTGATGTGGTGCTGGTGATCCCCTTCCAGAACCCT GAAGAGCAGGATGAAGGCTGGCTCATGGGCGTGAAGGAGCGACTGGAACCAGCACAAGGAGCTGGAG AAGTGCCGTGGCGTCTTCCCCGAGAACTTCACTGAGAGGGTCCCA<mark>TGA</mark>

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:

Sgfl-Mlul

Plasmid Map:



ACCN: NM_139349

Insert Size: 1428 bp

OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation:

This TrueClone is provided through our Custom Cloning Process that includes sub-cloning into OriGene's pCMV6 vector and full sequencing to provide a non-variant match to the expected reference without frameshifts, and is delivered as lyophilized plasmid DNA.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 139349.2</u>

RefSeq Size: 2332 bp
RefSeq ORF: 1428 bp
Locus ID: 274
UniProt ID: 000499

Cytogenetics: 2q14.3

MW: 51.7 kDa

Gene Summary:

This gene encodes several isoforms of a nucleocytoplasmic adaptor protein, one of which was initially identified as a MYC-interacting protein with features of a tumor suppressor. Isoforms that are expressed in the central nervous system may be involved in synaptic vesicle endocytosis and may interact with dynamin, synaptojanin, endophilin, and clathrin. Isoforms that are expressed in muscle and ubiquitously expressed isoforms localize to the cytoplasm and nucleus and activate a caspase-independent apoptotic process. Studies in mouse suggest that this gene plays an important role in cardiac muscle development. Alternate splicing of the gene results in several transcript variants encoding different isoforms. Aberrant splice variants expressed in tumor cell lines have also been described. [provided by RefSeq, Mar 2016]

Transcript Variant: This variant (7) lacks three in-frame exons in the coding region, compared to variant 1. Isoform 7, also called IIc2 and S1/R3-c, is shorter than isoform 1 and binds dynamin and synaptojanin.