

Product datasheet for **SC330083**

IL5RA (NM_001243099) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: IL5RA (NM_001243099) Human Untagged Clone
Tag: Tag Free
Symbol: IL5RA
Synonyms: CD125; CDw125; HSIL5R3; IL5R
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC330083 representing NM_001243099.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

ATGATCATCGTGGCGCATGTATTACTCATCCTTTTGGGGGCCACTGAGATACTGCAAGCTGACTTACTT
CCTGATGAAAAGATTTCACTTCTCCACCTGTCAATTTACCATTAAAGTTACTGGTTTGGCTCAAGTT
CTTTTACAATGGAACCAATCCTGATCAAGAGCAAAGGAATGTTAATCTAGAATATCAAGTAAAAATA
AACGCTCCAAAAGAAGATGACTATGAAACCAGAATCACTGAAAGCAAATGTGAACCATCTCCACAAA
GGCTTTTACAGCAAGTGTGCGGACCATCTGCAGAACGACCACTCACTACTGGCCAGCAGCTGGGCTTCT
GCTGAAC TTCATGCCCCACCAGGTCTCCTGGAACCTCAATTGTGAATTTAACTGCACCACAAACT
ACAGAAGACAATTATTCACGTTTAAAGGTATACCAAGTTTCCCTTCACTGCACCTGGCTTGTGGCACA
GATGCCCTGAGGACACGCAGTATTTTCTACTATAGGTATGGCTCTTGGACTGAAGAATGCCAAGAA
TACAGCAAAGACACACTGGGGAGAAATATCGCATGCTGGTTTCCAGGACTTTTATCCTCAGCAAAGGG
CGTGACTGGCTTGCAGTCTTGTAAACGGCTCCAGCAAGCACTCTGCTATCAGGCCCTTTGATCAGCTG
TTTGCCCTTACGCCATTGATCAAATAAATCCTCCACTGAATGTCACAGCAGAGATTGAAGGAACCTCGT
CTCTCTATCCAATGGGAGAAACCAGTGTCTGCTTTTCCAATCCATTGCTTTGATTATGAAGTAAAAATA
CACAATAACAAGGAATGGATATTTGCAGATAGAAAAATTGATGACCAATGCATTCTCAATAATTGAT
GATCTTTCTAAGTACGATGTTCAAGTGAGAGCAGCAGTGACTCCATGTGCAGAGAGGCAGGGCTCTGG
AGTGAGTGGAGCCAACCTATTTATGTGGGAAATGATGAACACAAGCCCTTGAGAGAGTGGTTTGTGATT
GTGATTATGGCAACCATCTGCTTCATCTTGTAAATCTCTCGCTTATCTGTAATAAAGCTGGGTCCA
GTGAGACGAAATTGAAGTCATCTGTTATAG
  
```

Restriction Sites: Sgfl-Mlul



[View online »](#)

Plasmid Map:


ACCN: NM_001243099

Insert Size: 1137 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001243099.1](#)

RefSeq Size: 5667 bp

RefSeq ORF:	1137 bp
Locus ID:	3568
UniProt ID:	Q01344
Cytogenetics:	3p26.2
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT signaling pathway
MW:	42.9 kDa

Gene Summary: The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported. [provided by RefSeq, Jul 2011]

Transcript Variant: This variant (7) differs in the 5' UTR and lacks an alternate coding exon compared to variant 1, that causes a frameshift. The resulting isoform (4) has a shorter and distinct C-terminus compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.