

## Product datasheet for SC333523

### Diazepam Binding Inhibitor (DBI) (NM\_001282633) Human Untagged Clone


#### Product data:

|                      |  |
|----------------------|--|
| Product Type:        | Expression Plasmids  |
| Product Name:        | Diazepam Binding Inhibitor (DBI) (NM_001282633) Human Untagged Clone                       |
| Tag:                 | Tag Free   |
| Symbol:              | DBI  |
| Synonyms:            | ACBD1; ACBP; CCK-RP; EP  |
| Vector:              | pCMV6-Entry (PS100001)   |
| Fully Sequenced ORF: | >SC333523 representing NM_001282633.<br>Blue=Insert sequence Red=Cloning site Green=Tag(s) |

```
ATGTGGGGCGACCTCTGGCTCCTCCCGCCTGCCTCTGCCAATCCGGGCACTGGGACAGAGGCTGAGTTT
GAGAAAGCTGCAGAGGAGGTTAGGCACCTTAAGACCAAGCCATCGGATGAGGAGATGCTGTTTCATCTAT
GGCCACTACAAACAAGCAACTGTGGGCGACATAAATACAGAACGGCCCGGGATGTTGGACTTCACGGGC
AAGCCAAGTGGGATGCCTGGAATGAGCTGAAAGGGACTTCCAAGGAAGATGCCATGAAAGCTTACATC
AACAAAGTAGAAGAGCTAAAGAAAAAATACGGGATATGA
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM\_001282633

Insert Size: 315 bp



[View online »](#)

|                               |  |
|-------------------------------|--|
| <b>OTI Disclaimer:</b>        | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).   |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>  |
| <b>RefSeq:</b>                | <a href="#">NM_001282633.1</a>   |
| <b>RefSeq Size:</b>           | 750 bp   |
| <b>RefSeq ORF:</b>            | 315 bp   |
| <b>Locus ID:</b>              | 1622   |
| <b>UniProt ID:</b>            | <a href="#">P07108</a>   |
| <b>Cytogenetics:</b>          | 2q14.2   |
| <b>Protein Families:</b>      | Druggable Genome   |
| <b>Protein Pathways:</b>      | PPAR signaling pathway   |
| <b>MW:</b>                    | 11.8 kDa   |
| <b>Gene Summary:</b>          | <p>This gene encodes diazepam binding inhibitor, a protein that is regulated by hormones and is involved in lipid metabolism and the displacement of beta-carbolines and benzodiazepines, which modulate signal transduction at type A gamma-aminobutyric acid receptors located in brain synapses. The protein is conserved from yeast to mammals, with the most highly conserved domain consisting of seven contiguous residues that constitute the hydrophobic binding site for medium- and long-chain acyl-Coenzyme A esters. Diazepam binding inhibitor is also known to mediate the feedback regulation of pancreatic secretion and the postprandial release of cholecystokinin, in addition to its role as a mediator in corticotropin-dependent adrenal steroidogenesis. Three pseudogenes located on chromosomes 6, 8 and 16 have been identified. Multiple transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]</p> <p>Transcript Variant: This variant (8) has an alternate splice site and an alternate exon in the 5' region, which results in an alternate translation start codon, compared to variant 5. The resulting isoform (1) has a shorter and distinct N-terminus, compared to isoform 5. Variants 1, 7, 8, 9, and 10 encode the same isoform 1.</p> |