

Product datasheet for **SC337276**

Hydroxysteroid (17 beta) Dehydrogenase 4 (HSD17B4) (NM_001292027) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Hydroxysteroid (17 beta) Dehydrogenase 4 (HSD17B4) (NM_001292027) Human Untagged Clone
Tag:	Tag Free
Symbol:	HSD17B4
Synonyms:	DBP; MFE-2; MFP-2; MPF-2; PRLTS1; SDR8C1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001292027, the custom clone sequence may differ by one or more nucleotides

```

ATGGAGAAGATCATTTCACAATGCAGATTCCTTTGTTTCAATGAATGATTTGGGAGGGGACTTCAAAGGAG
TTGGTAAAGGCTCCTTAGCTGCTGATAAGGTTGTTGAAGAAATAAGAAGGAGAGGTTGAAAAAGCAGTGGC
CAACTATGATTCAGTGGAAAGAAGGAGAGAAGGTTGTGAAGACAGCCCTGGATGCTTTTGGAAAGAATAGAT
GTTGTGGTCAACAATGCTGGAATTCTGAGGGATCGTTCCTTTGCTAGGATAAGTGATGAAGACTGGGATA
TAATCCACAGAGTTCATTTGCGGGTTCATTCCAAGTGACACGGGCAGCATGGGAACACATGAAGAAACA
GAAGTATGGAAGGATTATTATGACTTCATCAGCTTCAGGAATATATGGCAACTTTGGCCAGGCAATTAT
AGTGCTGCAAAGTTGGGTCTTCTGGCCTTGCAAATTCTTTGCAATTGAAGGCAGGAAAAGCAACATTC
ATTGTAACACCATTGCTCCTAATGCGGGATCACGGATGACTCAGACAGTTATGCCTGAAGATCTTGTGGA
AGCCCTGAAGCCAGAGTATGTGGCACCTCTTGCTTTGGCTTTGTCACGAGAGTTGTGAGGAGAATGGT
GGCTTGTGTTGAGTTGGAGCAGGATGGATTGAAAAATTACGCTGGGAGCGGACTCTTGGAGCTATTGTAA
GACAAAAGAATCACCCAATGACTCCTGAGGCAGTCAAGGCTAACTGGAAGAAGATCTGTGACTTTGAGAA
TGCCAGCAAGCCTCAGAGTATCCAAGAATCAACTGGCAGTATAATTGAAGTTCTGAGTAAAAATAGATTCA
GAAGGAGGAGTTTCAGCAATCATACTAGTCGTGCAACGCTCTACAGCAACATCAGGATTTGCTGGAGCTA
TTGGCCAGAAACTCCCTCCATTTTCTTATGCTTATACGGAAGCTGGAAGCTATTATGTATGCCCTTGGAGT
GGGAGCGTCAATCAAGGATCCAAAAGATTTGAAATTTATTTATGAAGGAAGTTCTGATTTCTCTGTTTG
CCCACCTTCGGAGTTATCATAGGTCAGAAATCTATGATGGGTGGAGGATTAGCAGAAAATCCTGGACTTT
CAATCAACTTTGCAAAGGTTCTTCATGGAGAGCAGTACTTAGAGTTATATAAACCACTTCCCAGAGCAGG
AAAATTTAAAATGTGAAGCAGTTGTTGCTGATGTCCTAGATAAAGGATCCGGTGTAGTGATTATTATGGAT
GTCTATTCTTATCTGAGAAGGAACTTATGCCACAATCAGTTCTCTCTTTCTGTTGGCTGGAG
GCTTTGGTGAAAAACGGACATCAGACAAAAGTCAAGGTAGCTGTAGCCATACCTAATAGACCTCCTGATGC
TGTACTTACAGATACCACCTCTTAAATCAGGCTGCTTTGTACCGCCTCAGTGGAGACTGGAATCCCTTA
CACATTGATCCTAACTTTGCTAGTCTAGCAGGTTTGGACAAGCCCATATTACATGGATTATGTACATTTG
GATTTTCTGCCAGGCGTGTGTTACAGCAGTTTGCAGATAATGATGTGTCAAGATTCAAGGCAATTAAGGC
TCGTTTTGCAAACCAGTATATCCAGGACAACTCTACAACTGAGATGTGGAAGGAAGGAAACAGAAAT
CATTTTCAAACCAAGGTCCAAGAACTGGAGACATTGTCATTTCAAATGCATATGTGGATCTTGCACCAA
CATCTGGTACTTCAGCTAAGACACCCTCTGAGGCGGGAAGCTTCAGAGTACCTTTGATTTTGGGAAAT
AGGACGCCGCTAAAGGATATTGGGCTGAGGTGGTGAAGAAAGTAAATGCTGTATTTGAGTGGCATATA
ACCAAAGCGGAAATATTGGGGCTAAGTGGACTTTGACCTGAAAAGTGGTTCTGAAAAGTGTACCAAG
GCCCTGCAAAAAGGTGCTGCTGATACAACAATCATACTTTCAGATGAAGATTTTCATGGAGGTGGTCTGGG
CAAGCTTGACCCTCAGAAGGCATTTTCTAGTGGCAGGCTGAAGGCCAGAGGGAACATCATGCTGAGCCAG
AACTTCAGATGATTCTTAAAGACTACGCCAAGCTCTGA
    
```

Restriction Sites: SgfI-MluI

ACCN: NM_001292027

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001292027.1, NP_001278956.1</u>
RefSeq Size:	2775 bp
RefSeq ORF:	2139 bp
Locus ID:	3295
UniProt ID:	<u>P51659</u>
Cytogenetics:	5q23.1
Protein Families:	Druggable Genome
Protein Pathways:	Metabolic pathways, Primary bile acid biosynthesis
Gene Summary:	<p>The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is present on chromosome 8. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014]</p> <p>Transcript Variant: This variant (4) has an alternate splice site in the 5' region, which causes translation initiation at a downstream AUG start codon, compared to variant 1. The resulting isoform (4) has a shorter N-terminus, compared to isoform 1.</p>