

Product datasheet for **SM1742PS**

B7-1 (CD80) Mouse Monoclonal Antibody [Clone ID: MEM-233]

Product data:

Product Type:	Primary Antibodies
Clone Name:	MEM-233
Applications:	FC, IP
Recommended Dilution:	Flow cytometry: 1-10 µg/ml. Immunoprecipitation.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Extracellular domain of human CD80 fused to human IgG1(Fc)
Specificity:	The antibody reacts with CD80 (B7-1), a 60 kDa single chain type I glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.
Formulation:	Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4 State: Purified State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein-A affinity chromatography; purity: > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.
Gene Name:	CD80 molecule
Database Link:	Entrez Gene 941 Human P33681




[View online »](#)

Background:

CD80 (B7-1) and CD86 (B7-2) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CTLA-4 (CD152). The both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, the both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.

Synonyms:

CD28LG, CD28LG1, LAB7, BB1, B7.1, B7-1

Product images:

Surface staining of RAJI human Burkitt lymphoma cell line with anti-human CD80 antibody (MEM-233) APC. Total viable cells were used for analysis.