

Product datasheet for SR415639

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sufu Mouse siRNA Oligo Duplex (Locus ID 24069)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: <u>NM 001025391, NM 015752</u>

UniProt ID: Q9Z0P7
Synonyms: Su(fu)

Components: Sufu (Mouse) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 24069)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml



Summary:

Negative regulator in the hedgehog/smoothened signaling pathway (PubMed:16155214, PubMed:16459298). Down-regulates GLI1-mediated transactivation of target genes (PubMed:11960000). Part of a corepressor complex that acts on DNA-bound GLI1 (PubMed:11960000). May also act by linking GLI1 to BTRC and thereby targeting GLI1 to degradation by the proteasome (By similarity). Sequesters GLI1, GLI2 and GLI3 in the cytoplasm, this effect is overcome by binding of STK36 to both SUFU and a GLI protein (PubMed:10531011, PubMed:16459298). Negative regulator of beta-catenin signaling (PubMed:11477086). Regulates the formation of either the repressor form (GLI3R) or the activator form (GLI3A) of the full-length form of GLI3 (GLI3FL) (PubMed:10531011, PubMed:20360384). GLI3FL is complexed with SUFU in the cytoplasm and is maintained in a neutral state (PubMed:10531011, PubMed:20360384). Without the Hh signal, the SUFU-GLI3 complex is recruited to cilia, leading to the efficient processing of GLI3FL into GLI3R (PubMed:10531011, PubMed:20360384). When Hh signaling is initiated, SUFU dissociates from GLI3FL and the latter translocates to the nucleus, where it is phosphorylated, destabilized, and converted to a transcriptional activator (GLI3A) (PubMed:10531011, PubMed:20360384). Required for normal embryonic development (PubMed:16155214, PubMed:16459298). Required for the proper formation of hair follicles and the control of epidermal differentiation (PubMed:16155214, PubMed:16459298, PubMed:23034632). [UniProtKB/Swiss-Prot Function]

Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).