

## **Product datasheet for TA317483**

## **UBC Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IF (1:10 - 1:50), IHC-P (10  $\mu$ g/ml), WB (1:1000)

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: UBC / Ubiquitin C antibody was raised against kLH conjugated synthetic peptide selected

from the C-terminal region of human Ubiquitin.

Formulation: PBS, 0.09% sodium azide

Purification: Protein G Column

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: ubiquitin C

Database Link: NP 066289

Entrez Gene 7316 Human

P0CG48

Synonyms: HMG20

**Protein Families:** Druggable Genome

**Protein Pathways:** PPAR signaling pathway


**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Anti-UBC / Ubiquitin C antibody IHC of human prostate. Immunohistochemistry of formalinfixed, paraffin-embedded tissue after heatinduced antigen retrieval. Antibody concentration 10 ug/ml.



Anti-UBC / Ubiquitin C antibody IHC of human testis. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody concentration 10 ug/ml.





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry;.