

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA321036

ACADS Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 200-1000 WB positive control: Human liver cancer tissue, rat kidney and human fetal liver tissue IHC: 25-100 Positive control: Human colon cancer Predicted cell location: Cytoplasm
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein corresponding to C terminal 250 amino acids of human acyl-CoA dehydrogenase, C-2 to C-3 short chain
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	44 kDa
Gene Name:	acyl-CoA dehydrogenase, C-2 to C-3 short chain
Database Link:	<u>NP_000008</u> <u>Entrez Gene 11409 MouseEntrez Gene 64304 RatEntrez Gene 35 Human</u> <u>P16219</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	ACADS Rabbit Polyclonal Antibody – TA321036
Background:	This gene encodes a a tetrameric mitochondrial flavoprotein, which is a member of the acyl- CoA dehydrogenase family. This enzyme catalyzes the initial step of the mitochondrial fatty acid beta-oxidation pathway. Mutations in this gene have been associated with Short Chain Acyl-CoA Dehydrogenase Deficiency. It is an autosomal recessive disorder resulting in acute acidosis and muscle weakness in infants, and a form of lipid-storage myopathy in adults.
Synonyms:	ACAD3; SCAD
Protein Families	Druggable Genome
Protein Pathway	s: Butanoate metabolism, Fatty acid metabolism, Metabolic pathways, Valine, leucine and isoleucine degradation

Product images:

×//___




Gel: 10%SDS-PAGE Lysate: 40 µg Lane 1-3: Human liver cancer tissue rat kidney tissue human fetal liver tissue Primary antibody: TA321036 (ACADS Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 90 seconds



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321036 (ACADS Antibody) at dilution 1/20 (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321036 (ACADS Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321036 (ACADS Antibody) at dilution 1/20 (Original magnification: ×200)

Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321036 (ACADS Antibody) at dilution 1/20, treated with fusion protein. (Original magnification: ×200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US