

Product datasheet for TA321459

VEGF Receptor 2 (KDR) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Rat brain tissue

IHC: 50-200

Positive control: Human colon cancer

Predicted cell location: Cytoplasm, Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 1163-1176 amino acids of human

kinase insert domain receptor (a type III receptor tyrosine kinase)

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 152 kDa

Gene Name: kinase insert domain receptor

Database Link: NP 002244

Entrez Gene 16542 MouseEntrez Gene 25589 RatEntrez Gene 3791 Human

P35968



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor; known as kinase insert domain receptor; is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation; survival; migration; tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors; including Rab GTPase; P2Y purine nucleotide receptor; integrin alphaVbeta3; T-cell protein tyrosine phosphatase; etc.. Mutations of this gene are implicated in infantile capillary hemangiomas.?

Synonyms: CD309; FLK1; VEGFR; VEGFR2

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, Endocytosis, Focal adhesion, VEGF signaling pathway

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: Rat brain tissue


Primary antibody: TA321459 (KDR Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321459 (KDR Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321459 (KDR Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)