

Product datasheet for TA336521

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Apolipoprotein A V (APOA5) Mouse Monoclonal Antibody [Clone ID: 1G5G9]

Product data:

Product Type: Primary Antibodies

Clone Name: 1G5G9

Applications: CyTOF-ready, ELISA, FC, IF, IHC, WB

Recommended Dilution: Western Blot: 0.5 ug/ml, Immunohistochemistry-Paraffin: 5 ug/ml, Flow Cytometry: 1 ug per

million cells, ELISA: 1:100-1:2000, Immunohistochemistry: 5 ug/ml, ICC/IF (Negative), CyTOF-

ready

Reactivity: Human Host: Mouse

Isotype: IgG1, kappa
Clonality: Monoclonal

Immunogen: Ni-NTA purified full-length recombinant human Apolipoprotein A5 expressed in E. Coli strain

BL21 (DE3). [UniProt# Q6Q788]

Formulation: PBS, pH 7.4, 0.1% Sodium Azide. Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

Concentration: lot specific

Purification: Protein G purified

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 41 kDa

Gene Name: apolipoprotein A5

Database Link: NP 443200

Entrez Gene 116519 Human

Q6Q788





Background:

Apolipoprotein A-V (APOA5) belongs to apolipoprotein A1/A4/E family and is a component of several lipoprotein fractions, including VLDL, HDL and chylomicrons. Expressed at high levels in plasma and liver, APOA5 exists as a secreted protein and interacts with GPIHBP1. APOA5 is an important determinant of plasma triglyceride levels by both behaving as a potent stimulator of apo-CII lipoprotein lipase/LPL TG hydrolysis and an inhibitor of hepatic VLDL-TG production rate (without affecting the VLDL-apoB production rate). APOA5 activates lecithin: cholesterol acyltransferase (LCAT) but does not enhance efflux of cholesterol from macrophages. Transgenic mice overexpressing human APOA5 gene exhibit one-third lower plasma TG levels than wild type, whereas APOA5 knockout mice possess four times as much plasma TG levels, with no effect on plasma cholesterol levels. Clinically, defects in APOA5 are a cause of susceptibility to familial hypertriglyceridemia (FHTR) and hyperlipoproteinemia type 5 (HLPP5).

Synonyms: APOAV; RAP3


Note: This Apolipoprotein A5 antibody is useful in ELISA, Immunohistochemistry paraffin embedded

sections and Western blot.

Protein Families: Druggable Genome, Secreted Protein


Protein Pathways: PPAR signaling pathway

Product images:




Flow Cytometry: Apolipoprotein A5 Antibody (1G5G9) TA336521 - Intracellular flow cytometric staining of 1 x 10^6 CHO (A) and HEK-293 (B) cells using Apolipoprotein A5 antibody (dark blue). Isotype control shown in orange. An antibody concentration of 1 ug/1x10^6 cells was used.






Western Blot: Apolipoprotein A5 Antibody (1G5G9) TA336521 - Analysis of ApoA5 in HepG2 lysates



Western Blot: Apolipoprotein A5 Antibody (1G5G9) TA336521 - Detection of Apo A5 in HepG2 cell lysate.



Western Blot: Apolipoprotein A5 Antibody (1G5G9) TA336521 - Analysis in human serum total protein (lane 2). Lane 1 is a MW standard.