

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA342263

Foxa2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB
Reactivity:	Mouse
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	The immunogen for anti-Foxa2 antibody: synthetic peptide corresponding to a region of Mouse. Synthetic peptide located within the following region: VDVEGTDYLNGDLGWSSSVSDSDERGSMQSLGSDEGYSSATVKRAKLQDG
Formulation:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose. Note that this product is shipped as lyophilized powder to China customers.
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	48 kDa
Gene Name:	forkhead box A2
Database Link:	<u>NP_034576</u> <u>Entrez Gene 15376 Mouse</u> <u>P35583</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Foxa2 Rabbit Polyclonal Antibody – TA342263

Background: Foxa2 is a transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Foxa2 is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. In embryonic development Foxa2 is required for notochord formation. Foxa2 is involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; Foxa1 and Foxa2 seem to have at least in part redundant roles. Foxa2 modulates the transcriptional activity of nuclear hormone receptors; inhibits AR-mediated transcription from the LCN5 promoter. Foxa2 binds to fibrinogen beta promoter and is involved in IL6-induced fibrinogen beta transcriptional activation. Foxa2 interacts with the cis-acting regulatory regions of these genes. Foxa2 is involved in glucose homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta-cells and glucose homeostasis. Foxa2 is involved in regulation of fat metabolism; activates transcriptional programs of lipid metabolism and ketogenesis at low insulin state.

Synonyms:HNF-3-beta; HNF-3B; HNF3B; MGC19807; TCF3BNote:Immunogen Sequence Homology: Mouse: 100%; Rat: 86%

Product images:



WB Suggested Anti-Foxa2 Antibody Titration: 0.2-1 ug/ml; Positive Control: Mouse Thymus

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US