

Product datasheet for TA345203

GTF2IRD1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:IHC, WBRecommended Dilution:WB, IHCReactivity:HumanHost:RabbitIsotype:IgG

Clonality: Polyclonal

Immunogen: The immunogen for anti-GTF2IRD1 antibody: synthetic peptide directed towards the N

terminal of human GTF2IRD1. Synthetic peptide located within the following region:

MALLGKRCDVPTNGCGPDRWNSAFTRKDEIITSLVSALDSMCSALSKLNA

Formulation: Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2%

sucrose.

Note that this product is shipped as lyophilized powder to China customers.

Purification: Protein A purified
Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 106 kDa

Gene Name: GTF2I repeat domain containing 1

Database Link: NP 057412

Entrez Gene 9569 Human

Q9UHL9



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: GTF2IRD1 contains five GTF2I-like repeats and each repeat possesses a potential helix-loop-

helix (HLH) motif. It may have the ability to interact with other HLH-proteins and function as a

transcription factor or as a positive transcriptional regulator under the control of

Retinoblastoma protein. GTF2IRD1 is related to Williams-Beuren syndrome, a multisystem developmental disorder. Western blots using three different antibodies against three unique regions of this protein target confirm the same apparent molecular weight in our tests. The protein encoded by this gene contains five GTF2I-like repeats and each repeat possesses a potential helix-loop-helix (HLH) motif. It may have the ability to interact with other HLHproteins and function as a transcription factor or as a positive transcriptional regulator under the control of Retinoblastoma protein. This gene is deleted in Williams-Beuren syndrome, a multisystem developmental disorder caused by deletion of multiple genes at 7q11.23.

Alternative splicing of this gene generates at least 2 transcript variants.

Synonyms: BEN; CREAM1; GTF3; hMusTRD1alpha1; MUSTRD1; RBAP2; WBS; WBSCR11; WBSCR12


Note: Immunogen Sequence Homology: Dog: 100%; Rat: 100%; Human: 100%; Pig: 93%; Horse: 93%;

Bovine: 93%; Rabbit: 86%; Guinea pig: 79%

Protein Families: Druggable Genome, Transcription Factors


Protein Pathways: Basal transcription factors

Product images:



WB Suggested Anti-GTF2IRD1 Antibody Titration: 2.5 ug/ml; ELISA Titer: 1: 312500; Positive Control: Transfected 293T





Human kidney