

## **Product datasheet for TA350894**

## **PIAS1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Human testis and human placenta tissue

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human PIAS1

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glyceroln

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 72 kDa

**Gene Name:** protein inhibitor of activated STAT 1

Database Link: NP 057250

Entrez Gene 8554 Human

075925

**Background:** This gene encodes a member of the mammalian PIAS [protein inhibitor of activated STAT-1

(signal transducer and activator of transcription-1)] family. This member contains a putative zinc-binding motif and a highly acidic region. It inhibits STAT1-mediated gene activation and the DNA binding activity, binds to Gu protein/RNA helicase II/DEAD box polypeptide 21, and

interacts with androgen receptor (AR). It functions in testis as a nuclear receptor transcriptional coregulator and may have a role in AR initiation and maintenance of

spermatogenesis.

Synonyms: DDXBP1; GBP; GU; RH-II; ZMIZ3



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Protein Families:** Druggable Genome, Transcription Factors

Protein Pathways: Jak-STAT signaling pathway, Pathways in cancer, Small cell lung cancer, Ubiquitin mediated

proteolysis

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane 1-2: Human testis tissue human placenta tissue

Primary antibody: TA350894 (PIAS1 Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 seconds