

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA355303**

## GCDFP 15 (PIP) Mouse Monoclonal Antibody [Clone ID: 23A3]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	23A3
Applications:	IHC
Recommended Dilution:	1:40
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2a
Clonality:	Monoclonal
Immunogen:	Recombinant prokaryotic protein corresponding to the excreted domain of the gross cystic disease fluid protein (15 kD) molecule
Specificity:	Human gross cystic disease fluid protein (15 kD)
Formulation:	Liquid tissue culture supernatant containing sodium azide as a preservative
Conjugation:	Unconjugated
Storage:	Store at 2-8°C
Stability:	12 months
Gene Name:	prolactin induced protein
Database Link:	<u>Entrez Gene 5304 Human</u> <u>P12273</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### GCDFP 15 (PIP) Mouse Monoclonal Antibody [Clone ID: 23A3] – TA355303

Background: Gross cystic disease of the breast is a benign premenopausal disorder in which cysts are a predominant pathological lesion. These cysts appear to be formed from excessive apocrine cystic secretions. This fluid is composed of several glycoproteins including a unique 15 kD monomer protein, GCDFP15. It has been reported that cytosolic analysis of normal tissue from all major organs has demonstrated GCDFP15 in apocrine epithelia, lacrimal, ceruminous and Moll's glands and in numerous serous cells of the submandibular, tracheal, bronchial, sublingual and minor salivary glands. Cytosol from breast carcinoma lesions are reported to contain GCDFP15 at a wide range of concentrations. The concentration is reported to be highest in more differentiated carcinomas and GCDFP15 shows only a few positive individual epithelial cells within lobules and small ducts in normal breast. Expression has also been reported in fibroadenomas within areas of apocrine metaplasia.

Synonyms:

GCDFP-15; GCDFP15; gp17; GPIP4; SABP

### **Product images:**



Invasive breast carcinoma: immunohistochemical staining for Gross Cystic Disease Fluid Protien-15. Gross Cystic Disease Fluid Protein-15: clone 23A3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US