

Product datasheet for TA373075

PD-L1 (CD274) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: 231, Raji, RAW264.7, NIH/3T3, HepG2, K562 cell lysates

Reactivity: Human, Mouse

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human CD274

pH7.4 PBS, 0.05% NaN3, 40% Glycerol Formulation:

Concentration: lot specific

Purification: Antigen affinity purification

Unconjugated Conjugation: Store at -20°C. Storage:

Stability: 1 year **Predicted Protein Size:** 33 kDa

Gene Name: CD274 molecule

Database Link: Entrez Gene 29126 Human

Q9NZQ7



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

Synonyms:

B7-H; B7-H1; B7H1; MGC142294; MGC142296; PD-L1; PDCD1L1; PDCD1LG1; PDL1

Product images:



Gel: 8%SDS-PAGE
Lysate: 40 µg
Lane 1-6: 231
Raji
RAW264.7
NIH/3T3
HepG2
K562 cell lysates
Primary antibody: TA373075 (CD274 Antibody) at dilution 1/600
Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution
Exposure time: 30 seconds