

Product datasheet for TA375932

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Eph receptor A4 (EPHA4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 887-986 of

human EPHA4 (NP_004429.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 104kDa/109kDa

Gene Name: EPH receptor A4

Database Link: Entrez Gene 2043 Human

P54764

Background: This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH


and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Multiple transcript variants encoding different isoforms have been found for this gene.




Synonyms:

HEK8; OTTHUMP00000164185; SEK; TYRO1

Product images:



Western blot analysis of extracts of various cell lines, using EPHA4 antibody (TA375932) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 3s.



Immunofluorescence analysis of C6 cells using EPHA4 antibody (TA375932) at dilution of 1:100. Blue: DAPI for nuclear staining.