

Product datasheet for TA375997

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Estrogen Related Receptor alpha (ESRRA) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-423 of

human ESRRA (NP_004442.3).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 45kDa

Gene Name: estrogen related receptor alpha

Database Link: Entrez Gene 2101 Human

P11474






Background:

The protein encoded by this gene is a nuclear receptor that is closely related to the estrogen receptor. This protein acts as a site-specific transcription regulator and has been also shown to interact with estrogen and the transcripton factor TFIIB by direct protein-protein contact. The binding and regulatory activities of this protein have been demonstrated in the regulation of a variety of genes including lactoferrin, osteopontin, medium-chain acyl coenzyme A dehydrogenase (MCAD) and thyroid hormone receptor genes. A processed pseudogene of ESRRA is located on chromosome 13q12.1. Alternatively spliced transcript variants have been found for this gene.

Synonyms:

ERR-alpha; ERR1; ERRa; ERRalpha; ESRL1; NR3B1; OTTHUMP00000198091

Product images:



Western blot analysis of extracts of various cell lines, using ESRRA antibody (TA375997) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST.