

Product datasheet for TA377430

Product data:

Product Type: Primary Antibodies

IL17RA Rabbit Polyclonal Antibody

Applications: WE

Reactivity: WB,1:1000 - 1:4000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 697-866 of

human IL17RA (NP_055154.3).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 92kDa/96kDa

Gene Name: interleukin 17 receptor A

Database Link: Entrez Gene 23765 Human

Q96F46

Background: Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It

is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to

interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript

variants encoding different isoforms.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

CD217; CDw217; hIL-17R; IL-17RA; IL17R; MGC10262

Product images:



Western blot analysis of extracts of various cell lines, using IL17RA antibody (TA377430) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 5s.