

## Product datasheet for TA500076BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### MAPK6 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1A9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A9
Applications: IHC, WB

**Reactivity:** WB: 1:1000, IHC 1;50 Human, Mouse, Rat

**Host:** Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Recombinant protein expressed in E.coli corresponding to amino acids 345-721 of human

ERK3

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 82.5 kDa

**Gene Name:** mitogen-activated protein kinase 6

Database Link: NP 002739

Entrez Gene 50772 MouseEntrez Gene 58840 RatEntrez Gene 5597 Human

016659

**Background:** ERK3 is a member of the Ser/Thr protein kinase family, and is most closely related to

mitogen-activated protein kinases (MAP kinases). MAP kinases also known as extracellular signal-regulated kinases (ERKs), are activated through protein phosphorylation cascades and act as integration points for multiple biochemical signals. This kinase is localized in the nucleus, and has been reported to be activated in fibroblasts upon treatment with serum or

phorbol esters







**Synonyms:** ERK3; HsT17250; p97MAPK; PRKM6

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAPK6 ([RC201597], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPK6. Positive lysates [LY400969] (100ug) and [LC400969] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded lung within the normal limits using anti-ERK3 ([TA000076]) mouse monoclonal antibody (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min).