

Product datasheet for TA500386AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TrkB (NTRK2) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2E1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2E1

Applications: IF, IHC, WB

Recommended Dilution: WB 1:1000, IHC 1:150, IF 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NTRK2 (NP_001007098) produced in

HEK293T cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 92 kDa

Gene Name: neurotrophic receptor tyrosine kinase 2

Database Link: NP 001007098

Entrez Gene 18212 MouseEntrez Gene 25054 RatEntrez Gene 4915 Human

016620

Background: This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This

kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself

and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation. Mutations in this gene have been associated with obesity and mood disorders. Alternate transcriptional splice variants encoding different isoforms have been

found for this gene






Synonyms: GP145-TrkB; trk-B; TRKB

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: MAPK signaling pathway, Neurotrophin signaling pathway


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NTRK2 (Cat# [RC205129], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NTRK2(Cat# [TA500386]). Positive lysates [LY400391] (100ug) and [LC400391] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-NTRK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500386])



Anti-NTRK2 mouse monoclonal antibody ([TA500386]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NTRK2 ([RC205129]).