

Product datasheet for **TA500724S**

ERAB (HSD17B10) Mouse Monoclonal Antibody [Clone ID: OTI11A2]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI11A2
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:50~100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human HSD17B10 (NP_004484) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.89 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	26.9 kDa
Gene Name:	hydroxysteroid 17-beta dehydrogenase 10
Database Link:	NP_004484 Entrez Gene 15108 Mouse Entrez Gene 63864 Rat Entrez Gene 3028 Human Q99714



[View online »](#)


Background: This gene encodes 3-hydroxyacyl-CoA dehydrogenase type II, a member of the short-chain dehydrogenase/reductase superfamily. The gene product is a mitochondrial protein that catalyzes the oxidation of a wide variety of fatty acids, alcohols, and steroids. The protein has been implicated in the development of Alzheimer's disease, and mutations in the gene are the cause of 2-methyl-3-hydroxybutyryl-CoA dehydrogenase deficiency (MHBD). Several alternatively spliced transcript variants have been identified, but the full-length nature of only two transcript variants has been determined.

Synonyms: 17b-HSD10; ABAD; CAMR; DUPXp11.22; ERAB; HADH2; HCD2; MHBD; MRPP2; MRX17; MRX31; MRXS10; SCHAD


Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Metabolic pathways, Valine, leucine and isoleucine degradation


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HSD17B10 ([RC201734], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSD17B10 ([TA500724]). Positive lysates [LY401426] (100ug) and [LC401426] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-HSD17B10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500724])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-HSD17B10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500724])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-HSD17B10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500724])




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-HSD17B10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500724])




Flow cytometric Analysis of Jurkat cells, using anti-HSD17B10 antibody ([TA500724]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of HeLa cells, using anti-HSD17B10 antibody ([TA500724]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Immunofluorescent staining of HepG2 cells using anti-HSD17B10 mouse monoclonal antibody ([TA500724]).



Anti-HSD17B10 mouse monoclonal antibody ([TA500724]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HSD17B10 ([RC201734]).