

Product datasheet for **TA500886S**

XRCC1 Mouse Monoclonal Antibody [Clone ID: OTI2D8]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI2D8
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human XRCC1 (NP_006288) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.57 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	69.3 kDa
Gene Name:	X-ray repair cross complementing 1
Database Link:	NP_006288 Entrez Gene 22594 Mouse Entrez Gene 84495 Rat Entrez Gene 7515 Human P18887
Background:	The protein encoded by this gene is involved in the efficient repair of DNA single-strand breaks formed by exposure to ionizing radiation and alkylating agents. This protein interacts with DNA ligase III, polymerase beta and poly (ADP-ribose) polymerase to participate in the base excision repair pathway. It may play a role in DNA processing during meiosis and recombination in germ cells. A rare microsatellite polymorphism in this gene is associated with cancer in patients of varying radiosensitivity. [provided by RefSeq]




[View online »](#)

Synonyms: RCC
Protein Families: Druggable Genome
Protein Pathways: Base excision repair


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY XRCC1 ([RC204952], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-XRCC1 (1:1000).




Western blot analysis of extracts (10ug) from 4 different cell lines by using anti-XRCC1 monoclonal antibody at 1:200 dilution.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])



Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-XRCC1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500886])



HEK293T cells transfected with either [RC204952] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-XRCC1 antibody ([TA500886]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-XRCC1 antibody ([TA500886]), (Red), compared to a nonspecific negative control antibody, (Blue).