

Product datasheet for TA501421BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MIOX Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4G7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4G7

Applications: FC, IF, IHC, WB

Recommended Dilution: WB: 1:200 - 1:1000, IHC 1:50, IF 1:100, FLOW 1:100

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MIOX (NP_060054) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 32.8 kDa

Gene Name: myo-inositol oxygenase

Database Link: NP 060054

Entrez Gene 55586 Human

Q9UGB7


Synonyms: ALDRL6

Protein Pathways: Ascorbate and aldarate metabolism, Inositol phosphate metabolism







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MIOX (Cat# [RC210070], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MIOX(Cat# [TA501421]). Positive lysates [LY413679] (100ug) and [LC413679] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MIOX monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])




Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])




Immunohistochemical staining of paraffinembedded Human thyroid tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])







Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-MIOX mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA501421])



Anti-MIOX mouse monoclonal antibody ([TA501421]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY MIOX ([RC210070]).



HEK293T cells transfected with either [RC210070] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-MIOX antibody ([TA501421]), and then analyzed by flow cytometry.