

## Product datasheet for **TA502677BM**

### ATP6V1B1 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1H6]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI1H6
Applications:	FC, IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human ATP6V1B1 (NP_001683) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	56.7 kDa
Gene Name:	ATPase H <sup>+</sup> transporting V1 subunit B1
Database Link:	<a href="#">NP_001683</a> <a href="#">Entrez Gene 110935 Mouse</a> <a href="#">Entrez Gene 312488 Rat</a> <a href="#">Entrez Gene 525 Human</a> <a href="#">P15313</a>



[View online »](#)



**Background:**

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of two V1 domain B subunit isoforms and is found in the kidney. Mutations in this gene cause distal renal tubular acidosis associated with sensorineural deafness. [provided by RefSeq]

**Synonyms:**


ATP6B1; RTA1B; VATB; VMA2; VPP3

**Protein Families:**

Druggable Genome


**Protein Pathways:**

Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection


**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ATP6V1B1 (Cat# [RC209462], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ATP6V1B1 (Cat# [TA502677]). Positive lysates [LY400635] (100ug) and [LC400635] (20ug) can be purchased separately from OriGene.






Anti-ATP6V1B1 mouse monoclonal antibody ([TA502677]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ATP6V1B1 ([RC209462]).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-ATP6V1B1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502677])




Immunohistochemical staining of paraffin-embedded Human Kidney tissue within the normal limits using anti-ATP6V1B1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502677])






HEK293T cells transfected with either [RC209462] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ATP6V1B1 antibody ([TA502677]), and then analyzed by flow cytometry.



Flow cytometric Analysis of Jurkat cells, using anti-ATP6V1B1 antibody ([TA502677]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).



Flow cytometric Analysis of Hela cells, using anti-ATP6V1B1 antibody ([TA502677]), (Red), compared to a nonspecific negative control antibody (TA50011), (Blue).