

Product datasheet for TA503913AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Adiponectin (ADIPOQ) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7B11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7B11
Applications: FC, IF, WB

Recommended Dilution: IF 1:100, FLOW 1:100, WB 1:200

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 19-244 of human

ADIPOQ(NP_004788) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 24.9 kDa

Gene Name: adiponectin, C1Q and collagen domain containing

Database Link: NP 004788

Entrez Gene 9370 Human

Q15848

Background: This gene is expressed in adipose tissue exclusively. It encodes a protein with similarity to

collagens X and VIII and complement factor C1q. The encoded protein circulates in the plasma and is involved with metabolic and hormonal processes. Mutations in this gene are associated with adiponectin deficiency. Multiple alternatively spliced variants, encoding the

same protein, have been identified. [provided by RefSeq, Apr 2010]




Adiponectin (ADIPOQ) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI7B11] – TA503913AM

Synonyms: ACDC; ACRP30; ADIPQTL1; ADPN; APM-1; APM1; GBP28


Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Adipocytokine signaling pathway, PPAR signaling pathway, Type II diabetes mellitus

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ADIPOQ ([RC215161], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ADIPOQ. Positive lysates [LY401510] (100ug) and [LC401510] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 1 cell line by using anti-ADIPOQ monoclonal antibody at 1:200.





Anti-ADIPOQ mouse monoclonal antibody ([TA503913]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ADIPOQ ([RC215161]).



HEK293T cells transfected with either [RC215161] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ADIPOQ antibody ([TA503913]), and then analyzed by flow cytometry.