

# **Product datasheet for TA503933**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## N Cadherin (CDH2) Mouse Monoclonal Antibody [Clone ID: OTI2G7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G7

**Applications:** FC, IHC, WB

**Recommended Dilution:** WB 1:500~2000, IHC 1:150, FLOW 1:100

Reactivity: Human, Dog, Monkey, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CDH2(NP\_001783) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.77 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 97.2 kDa

Gene Name: cadherin 2

Database Link: NP 001783

Entrez Gene 12558 MouseEntrez Gene 83501 RatEntrez Gene 480169 DogEntrez Gene 711526

MonkeyEntrez Gene 1000 Human

P19022





**Background:** This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a

calcium dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. The protein functions during gastrulation and is required for establishment of left-right asymmetry. At certain central nervous system synapses, presynaptic to postsynaptic adhesion is mediated at


least in part by this gene product. [provided by RefSeq]

Synonyms: CD325; CDHN; CDw325; NCAD


**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** Arrhythmogenic right ventricular cardiomyopathy (ARVC), Cell adhesion molecules (CAMs)

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CDH2 (Cat# [RC207170], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDH2 (Cat# TA503933). Positive lysates [LY419743] (100ug) and [LC419743] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CDH2 monoclonal antibody.






Western blot analysis of extracts (35ug) from 6 cell lines by using anti-CDH2 monoclonal antibody (1:500).




Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and CDH2-Knockout 293T cells (KO, Cat# [LC840285]) were separated by SDS-PAGE and immunoblotted with anti-CDH2 monoclonal antibody TA503933, (1:500). Then the blotted membrane was stripped and reprobed with antib-actin antibody ([TA811000]) as a loading control.




Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-CDH2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503933) (1:150)






Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-CDH2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503933) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-CDH2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503933) (1:150)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-CDH2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503933) (1:150)









Flow cytometric Analysis of Hela cells, using anti-CDH2 antibody (TA503933), (Red), compared to a nonspecific negative control antibody, (Blue).