

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA505579

NDOR1 Mouse Monoclonal Antibody [Clone ID: OTI2G4]

Product data:

| Product Type: | Primary Antibodies |
|-------------------------|--|
| Clone Name: | OTI2G4 |
| Applications: | IF, IHC, WB |
| Recommended Dilution: | WB 1:200 - 1:500, IHC 1:150, IF 1:100 |
| Reactivity: | Human |
| Host: | Mouse |
| lsotype: | IgG2a |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human NDOR1(NP_055249) produced in HEK293T cell. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 1 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 66.6 kDa |
| Gene Name: | NADPH dependent diflavin oxidoreductase 1 |
| Database Link: | <u>NP_055249</u> <u>Entrez Gene 27158 Human</u> <u>Q9UHB4</u> |
| Background: | This gene encodes an NADPH-dependent diflavin reductase that contains both flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) binding domains. The encoded protein catalyzes the transfer of electrons from NADPH through FAD and FMN cofactors to potential redox partners. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2012] |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Synonyms:


bA350O14.9; NR1

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NDOR1 ([RC204845], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NDOR1. Positive lysates [LY415283] (100ug) and [LC415283] (20ug) can be purchased separately from OriGene.

Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-NDOR1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505579)



Anti-NDOR1 mouse monoclonal antibody (TA505579) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NDOR1 ([RC204845]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunofluorescent staining of HeLa cells using anti-NDOR1 mouse monoclonal antibody (TA505579) at 1:100 dilution.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US