

Product datasheet for TA506764AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BID Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2A2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A2
Applications: IHC, WB

Recommended Dilution: WB 1:4000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BID(NP_001187) produced in HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21.8 kDa

Gene Name: BH3 interacting domain death agonist

Database Link: NP 001187

Entrez Gene 637 Human

P55957

Synonyms: FP497

Protein Families: Druggable Genome


Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Natural killer cell

mediated cytotoxicity, p53 signaling pathway, Pathways in cancer, Viral myocarditis







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BID ([RC207261], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BID. Positive lysates [LY420074] (100ug) and [LC420074] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-BID mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506764])



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-BID mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA506764])