

Product datasheet for TA801300S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

NM23A (NME1) Mouse Monoclonal Antibody [Clone ID: OTI18C12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI18C12

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NME1 (NP_937818) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 19.5 kDa

Gene Name: NME/NM23 nucleoside diphosphate kinase 1

Database Link: NP 937818

Entrez Gene 18102 MouseEntrez Gene 191575 RatEntrez Gene 4830 Human

P15531



NM23A (NME1) Mouse Monoclonal Antibody [Clone ID: OTI18C12] - TA801300S

Background: This gene (NME1) was identified because of its reduced mRNA transcript levels in highly

metastatic cells. Nucleoside diphosphate kinase (NDK) exists as a hexamer composed of 'A' (encoded by this gene) and 'B' (encoded by NME2) isoforms. Mutations in this gene have been identified in aggressive neuroblastomas. Two transcript variants encoding different isoforms

have been found for this gene. Co-transcription of this gene and the neighboring

downstream gene (NME2) generates naturally-occurring transcripts (NME1-NME2), which encodes a fusion protein comprised of sequence sharing identity with each individual gene


product. [provided by RefSeq, Jul 2008]

Synonyms: AWD; GAAD; NB; NBS; NDKA; NDPK-A; NDPKA; NM23; NM23-H1

Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: Metabolic pathways, Purine metabolism, Pyrimidine metabolism

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NME1 ([RC220517], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NME1. Positive lysates [LY404982] (100ug) and [LC404982] (20ug) can be purchased separately from OriGene.