

Product datasheet for TA801918

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

USP44 Mouse Monoclonal Antibody [Clone ID: OTI1F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1F9
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 420-712 of human

USP44 (NP_115523) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 81 kDa

Gene Name: ubiquitin specific peptidase 44

Database Link: NP 115523

Entrez Gene 84101 Human

O9H0E7

Background: Modification of cellular proteins by ubiquitin is an essential regulatory mechanism controlled

by the coordinated action of multiple ubiquitin-conjugating and deubiquitinating enzymes. USP44 belongs to a large family of cysteine proteases that function as deubiquitinating enzymes (Quesada et al., 2004 [PubMed 14715245]). [supplied by OMIM, Mar 2008]


Synonyms: DKFZp434D0127; FLJ14528






Protein Families: Druggable Genome, Protease


Product images:



E.coli lysate (left lane) and E.coli lysate expressing human recombinant protein fragment corresponding to amino acids 420-712 of human USP44 (NP_115523) were separated by SDS-PAGE and immunoblotted with anti-USP44.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)






Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Human Ovary tissue within the normal limits using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)




Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)





Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-USP44 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA801918)