

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA802719S

## Ornithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI7B6]


## **Product data:**

Product Type:	Primary Antibodies	
Clone Name:	OTI7B6	
Applications:	IHC, WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 33-354 of human OTC (NP_000522)produced in SF9 cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	36.1 kDa	
Gene Name:	ornithine carbamoyltransferase	
Database Link:	<u>NP_000522</u> <u>Entrez Gene 18416 MouseEntrez Gene 25611 RatEntrez Gene 5009 Human</u> <u>P00480</u>	
Background:	This nuclear gene encodes a mitochondrial matrix enzyme. Missense, nonsense, and frameshift mutations in this enzyme lead to ornithine transcarbamylase deficiency, which causes hyperammonemia. Since the gene for this enzyme maps close to that for Duchenne muscular dystrophy, it may play a role in that disease also. [provided by RefSeq, Jul 2008]	
Synonyms:	OCTD; OTCD	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Ornithine Carbamoyltransfera TA802719S	rnithine Carbamoyltransferase (OTC) Mouse Monoclonal Antibody [Clone ID: OTI7B6] – A802719S	
Protein Familie	s: Druggable Genome		
Protein Pathwa	<b>ys:</b> Arginine and proline me	Arginine and proline metabolism, Metabolic pathways	
Product ima	ges:		
	$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	HEK293T cells were transfected with the pCMV6- ENTRY control (Left lane) or pCMV6-ENTRY OTC ([RC214662], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-OTC.	



15

10

Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-OTC mouse monoclonal antibody. ([TA802719]) Dilution: 1:150

Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-OTC mouse monoclonal antibody. ([TA802719]) Dilution: 1:150

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US