

## **Product datasheet for TA803506S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MSMB Mouse Monoclonal Antibody [Clone ID: OTI2E7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2E7

Applications: IHC, WB

Recommended Dilution: WB 1:1000

Reactivity: Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Formulation:** Purified prostate specific antigen (MSMB) from human seminal fluid PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 10.7 kDa

**Gene Name:** microseminoprotein, beta-

Database Link: NP 002434

Entrez Gene 4477 Human

P08118

**Background:** The protein encoded by this gene is a member of the immunoglobulin binding factor family.


It is synthesized by the epithelial cells of the prostate gland and secreted into the seminal plasma. This protein has inhibin-like activity. It may have a role as an autocrine paracrine factor in uterine, breast and other female reproductive tissues. The expression of the encoded protein is found to be decreased in prostate cancer. Two alternatively spliced transcript variants encoding different isoforms are described for this gene. The use of alternate polyadenylation sites has been found for this gene. [provided by RefSeq, Jul 2008]




Synonyms: HPC13; IGBF; MSP; MSPB; PN44; PRPS; PSP-94; PSP57; PSP94

**Protein Families:** Secreted Protein, Transmembrane

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MSMB ([RC202704], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MSMB. Positive lysates [LY400872] (100ug) and [LC400872] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-MSMB mouse monoclonal antibody. ([TA803506]) Dilution: 1:150