

Product datasheet for TA803667

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MDA5 (IFIH1) Mouse Monoclonal Antibody [Clone ID: OTI8C10]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: OTI8C10

Applications: WB

Recommended Dilution: WB 1:100
Reactivity: Human
Host: Mouse

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IFIH1 (NP_071451) produced in HEK293T

cell

lgG1

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 116.5 kDa

Gene Name: interferon induced with helicase C domain 1

Database Link: NP 071451

Entrez Gene 64135 Human

O9BYX4




Background:

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus insulin-dependent type 19. [provided by RefSeq, Jul 2012]

Synonyms: AGS7; Hlcd; IDDM19; MDA-5; MDA5; RLR-2

Protein Pathways: RIG-I-like receptor signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IFIH1 (Cat# [RC215661], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFIH1(Cat# TA803667). Positive lysates [LY411723] (100ug) and [LC411723] (20ug) can be purchased separately from OriGene.