

Product datasheet for **TA803782AM**

Fibronectin (FN1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3F9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3F9
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 32-307 of human FN1 (NP_473375) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	fibronectin 1
Database Link:	NP_473375 Entrez Gene 14268 Mouse Entrez Gene 25661 Rat Entrez Gene 2335 Human P02751
Background:	This gene encodes fibronectin, a glycoprotein present in a soluble dimeric form in plasma, and in a dimeric or multimeric form at the cell surface and in extracellular matrix. Fibronectin is involved in cell adhesion and migration processes including embryogenesis, wound healing, blood coagulation, host defense, and metastasis. The gene has three regions subject to alternative splicing, with the potential to produce 20 different transcript variants. However, the full-length nature of some variants has not been determined. [provided by RefSeq, Jul 2008]




[View online »](#)

Synonyms: CIG; ED-B; FINC; FN; FNZ; GFND; GFND2; LETS; MSF


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Protein Pathways: ECM-receptor interaction, Focal adhesion, Pathways in cancer, Regulation of actin cytoskeleton, Small cell lung cancer


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FN1 ([RC212860], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FN1. Positive lysates [LY409278] (100ug) and [LC409278] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Human liver tissue within the normal limits using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Human lung tissue within the normal limits using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-FN1 mouse monoclonal antibody. ([TA803782]) Dilution: 1:150



Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150



Immunohistochemical staining of paraffin-embedded Carcinoma of Human bladder tissue using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-FN1 mouse monoclonal antibody. (TA803782) Dilution: 1:150