

# **Product datasheet for TA805286**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## COX2 (PTGS2) Mouse Monoclonal Antibody [Clone ID: OTI5F12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5F12
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 18-260 of human PTGS2

(NP\_000954) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 67.2 kDa

**Gene Name:** prostaglandin-endoperoxide synthase 2

Database Link: NP 000954

Entrez Gene 19225 MouseEntrez Gene 29527 RatEntrez Gene 5743 Human

P35354

### COX2 (PTGS2) Mouse Monoclonal Antibody [Clone ID: OTI5F12] - TA805286

**Background:** Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key

enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb

2009]


**Synonyms:** COX-2; COX2; GRIPGHS; hCox-2; PGG/HS; PGHS-2; PHS-2

**Protein Families:** Druggable Genome


**Protein Pathways:** Arachidonic acid metabolism, Pathways in cancer, Small cell lung cancer, VEGF signaling

pathway

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PTGS2 (Cat# [RC202245], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PTGS2(Cat# TA805286).



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-PTGS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA805286)





Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-PTGS2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA805286)