

### Product datasheet for TA805461AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TLK2 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2F9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F9
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-296 of human TLK2

(NP\_006843) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 85.3 kDa

**Gene Name:** tousled like kinase 2

Database Link: NP 006843

Entrez Gene 24086 MouseEntrez Gene 303592 RatEntrez Gene 11011 Human

**086UE8** 


Synonyms: HsHPK; PKU-ALPHA

**Protein Families:** Druggable Genome, Protein Kinase






# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TLK2 ([RC218935], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TLK2. Positive lysates [LY416375] (100ug) and [LC416375] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-TLK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA805461])





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-TLK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA805461])