

Product datasheet for TA805507BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TLK2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3E10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3E10

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-296 of human

TLK2(NP_006843) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 85.3 kDa

Gene Name: tousled like kinase 2

Database Link: NP 006843

Entrez Gene 24086 MouseEntrez Gene 303592 RatEntrez Gene 11011 Human

Q86UE8

Background: This gene encodes a nuclear serine/threonine kinase that was first identified in Arabidopsis.

The encoded protein is thought to function in the regulation of chromatin assembly in the S phase of the cell cycle by regulating the levels of a histone H3/H4 chaperone. This protein is

associated with double-strand break repair of DNA damage caused by radiation.

Pseudogenes of this gene are present on chromosomes 10 and 17. Alternate splicing results

rseddogenes of this gene are present of chromosomes to and 17. Alternate splicing results

in multiple transcript variants. [provided by RefSeq, Sep 2013]







Synonyms: HsHPK; PKU-ALPHA

Protein Families: Druggable Genome, Protein Kinase

Product images:



Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-TLK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA805507]) (1:150)



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-TLK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA805507]) (1:150)