

## **Product datasheet for TA805600**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Natriuretic Peptide Receptor A (NPR1) Mouse Monoclonal Antibody [Clone ID: OTI7H7]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI7H7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 746-1006 of human

NPR1 (NP\_000897) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 118.7 kDa

**Gene Name:** natriuretic peptide receptor 1

Database Link: NP 000897

Entrez Gene 18160 MouseEntrez Gene 24603 RatEntrez Gene 4881 Human

P16066



# Natriuretic Peptide Receptor A (NPR1) Mouse Monoclonal Antibody [Clone ID: OTI7H7] – TA805600

Background:


Guanylyl cyclases, catalyzing the production of cGMP from GTP, are classified as soluble and membrane forms (Garbers and Lowe, 1994 [PubMed 7982997]). The membrane guanylyl cyclases, often termed guanylyl cyclases A through F, form a family of cell-surface receptors with a similar topographic structure: an extracellular ligand-binding domain, a single membrane-spanning domain, and an intracellular region that contains a protein kinase-like domain and a cyclase catalytic domain. GC-A and GC-B function as receptors for natriuretic peptides; they are also referred to as atrial natriuretic peptide receptor A (NPR1) and type B (NPR2; MIM 108961). Also see NPR3 (MIM 108962), which encodes a protein with only the ligand-binding transmembrane and 37-amino acid cytoplasmic domains. NPR1 is a membrane-bound guanylate cyclase that serves as the receptor for both atrial and brain natriuretic peptides (ANP (MIM 108780) and BNP (MIM 600295), respectively). [supplied by OMIM, May 2009]

Synonyms: ANPa; ANPRA; GUC2A; GUCY2A; NPRA

Protein Families: Druggable Genome, Protein Kinase

**Protein Pathways:** Purine metabolism, Vascular smooth muscle contraction

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NPR1 (Cat# [RC209267], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NPR1(Cat# TA805600). Positive lysates [LY424461] (100ug) and [LC424461] (20ug) can be purchased separately from OriGene.