

Product datasheet for TA806459BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Retinoic Acid Receptor alpha (RARA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7B8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7B8

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 52-299 of human

RARA(NP 000955) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50.6 kDa

Gene Name: retinoic acid receptor alpha

Database Link: NP 000955

Entrez Gene 19401 MouseEntrez Gene 24705 RatEntrez Gene 5914 Human

P10276





Retinoic Acid Receptor alpha (RARA) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7B8] – TA806459BM

Background: This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid

receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoeisis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants


have been found for this locus. [provided by RefSeq, Sep 2010]

Synonyms: NR1B1; RAR

Protein Families: Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

Protein Pathways: Acute myeloid leukemia, Pathways in cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RARA ([RC200430], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RARA. Positive lysates [LY400350] (100ug) and [LC400350] (20ug) can be purchased separately from OriGene.