

Product datasheet for TA811947

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TOMM40 Mouse Monoclonal Antibody [Clone ID: OTI9A3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9A3
Applications: IHC, WB

Reactivity: WB 1:500, IHC 1:500 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TOMM40 (NP_006105) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 37.7 kDa

Gene Name: translocase of outer mitochondrial membrane 40

Database Link: NP 006105

Entrez Gene 53333 MouseEntrez Gene 308416 RatEntrez Gene 10452 Human

096008

Background: The protein encoded by this gene is localized in the outer membrane of the mitochondria. It

is the channel-forming subunit of the translocase of the mitochondrial outer membrane (TOM) complex that is essential for import of protein precursors into mitochondria.

Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq,

Aug 2015]

Synonyms: C19orf1; D19S1177E; PER-EC1; PEREC1; TOM40







Protein Families: Druggable Genome, Ion Channels: Other

Protein Pathways: Amyotrophic lateral sclerosis (ALS)

Product images:




Western blot analysis of extracts (35ug) from 2 cell lines and human liver tissue lysate by using anti-TOMM40 monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-TOMM40 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA811947) (1:500)





Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-TOMM40 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA811947) (1:500)