

Product datasheet for TA813088BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

B7-2 (CD86) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI11E3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI11E3
Applications: FC, WB

Recommended Dilution: WB 1:500~1000, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD86 (NP_787058) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 38.1 kDa

Gene Name: CD86 molecule

Database Link: NP 787058

Entrez Gene 942 Human

P42081





Background: This gene encodes a type I membrane protein that is a member of the immunoglobulin

superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms. [provided by RefSeq, May 2011]

Synonyms: B7-2; B7.2; B70; CD28LG2; LAB72


Protein Families: Druggable Genome, Transcription Factors, Transmembrane

Protein Pathways: Allograft rejection, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-


versus-host disease, Systemic lupus erythematosus, Toll-like receptor signaling pathway,

Type I diabetes mellitus, Viral myocarditis


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD86 ([RC217341], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD86 (1:1000).



Western blot analysis of extracts (35ug) from 3 cell lines lysates by using anti-CD86 monoclonal antibody (1:500).




Flow cytometric analysis of living 293T cells transfected with CD86 overexpression plasmid ([RC217341]), Red) using anti-CD86 antibody ([TA813088]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).



Flow cytometric analysis of living Raji cells, using anti-CD86 antibody ([TA813088], Red), compared to an isotype control (green) (1:100).





Flow cytometric analysis of living Ramos cells, using anti-CD86 antibody ([TA813088], Red), compared to an isotype control (green) (1:100).