

Product datasheet for TA813845

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD38 Mouse Monoclonal Antibody [Clone ID: OTI11C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI11C3

Applications: FC

Recommended Dilution: FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD38 (NP_001766) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 34.3 kDa

Gene Name: CD38 molecule

Database Link: NP 001766

Entrez Gene 952 Human

P28907

Background: Synthesizes the second messagers cyclic ADP-ribose and nicotinate-adenine dinucleotide

phosphate, the former a second messenger for glucose-induced insulin secretion. Also has cADPr hydrolase activity. Also moonlights as a receptor in cells of the immune system.

[UniProtKB/Swiss-Prot Function]





Synonyms: ADPRC 1; ADPRC1


Protein Families: Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem

cells, Transmembrane


Protein Pathways: Calcium signaling pathway, Hematopoietic cell lineage, Metabolic pathways, Nicotinate and

nicotinamide metabolism

Product images:




Flow cytometric analysis of living 293T cells transfected with CD38 overexpression plasmid ([RC203179]), Red)/empty vector ([PS100001], Blue) using anti-CD38 antibody (TA813845). Cells incubated with a non-specific antibody (Green) were used as isotype control. [] [] [] [] [] []




Flow cytometric analysis of living Raji cells, using anti-CD38 antibody(TA813845, Red), compared to an isotype control (green).(1:100)






Flow cytometric analysis of living K562 cells, using anti-CD38 antibody(TA813845, Red), compared to an isotype control (green), and a PBS control (blue).(1:100)



Flow cytometric analysis of living RPMI-8226 cells, using anti-CD38 antibody(TA813845, Red), compared to an isotype control (green), and a PBS control (blue).(1:100)



Flow cytometric analysis of living OPM2 cells, using anti-CD38 antibody(TA813845, Red), compared to an isotype control (green), and a PBS control (blue).(1:100)