

## Product datasheet for TA814158BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HSD17B13 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5C4]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5C4
Applications: IHC, WB

Recommended Dilution: WB 1:500-1:1000, IHC 1:200

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 20-300 of human

HSD17B13 produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 33.7 kDa

**Gene Name:** hydroxysteroid (17-beta) dehydrogenase 13

Database Link: NP 835236

Entrez Gene 345275 Human

O7Z5P4


Synonyms: HMFN0376; NIIL497; SCDR9; SDR16C3

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane





# **Product images:**




Western blot analysis of overexpressed human HSD17B13 lysate (5ug [LY405941], LEFT) and purified recombinant protein (0.02ug [TP313132], RIGHT) by using anti-HSD17B13 antibody ([TA814158]).




HEK293T cells were transfected with 10 different plasmids of human HSD cDNA (HSD11B1, HSD17B2, HSD17B3, HSD17B4, HSD17B6, HSD17B7, HSD17B8, HSD17B10, HSD17B11, HSD17B13) for 48 hrs and lysed. Equivalent amounts of cell lysates (10ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSD17B13 mouse monoclonal antibody. HSD17B13 was positive, while all others were negative. ([TA814158], 10500)





Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-HSD17B13 mouse monoclonal antibody, [TA814158] (1:200). \*Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-HSD17B13 mouse monoclonal antibody, [TA814158] (1:200). \*Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3 min.