

Product datasheet for TA815233S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL17E (IL25) Mouse Monoclonal Antibody [Clone ID: OTI12G2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI12G2

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IL17E (NP_073626) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Shipped at -20°C or with ice packs, Upon delivery store at -20°C. Dilute in PBS(pH7.3) if

necessary. Stable for 12 months from date of receipt. Avoid repeated freeze-thaws.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 20.3 kDa

Gene Name: interleukin 25

Database Link: NP 073626

Entrez Gene 64806 Human

Q9H293

Background: The protein encoded by this gene is a cytokine that shares sequence similarity with

interleukin 17. This cytokine can induce NF-kappaB activation, and stimulate the production of interleukin 8. Both this cytokine and interleukin 17B are ligands for the cytokine receptor

IL17BR. Studies of a similar gene in mice suggest that this cytokine may be a pro-

inflammatory cytokine favoring the Th2-type immune response. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Mar 2010]





Synonyms: IL17E

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction

Product images:




Figure A, Western blot analysis of overexpressed lysates(15ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], lane 1), human IL17E plasmid ([RC210141], lane 2), mouse IL17E plasmid ([MR224044], lane 3) using anti-IL17E antibody [TA815233] (1:2000). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)